

Biosynthesis of the Methanogenic Cofactors

ROBERT H. WHITE

*Department of Biochemistry, Virginia Polytechnic Institute
and State University, Blacksburg, Virginia 24061*

- I. Introduction
- II. Biosynthesis of Methanopterin
- III. Biosynthesis of Coenzyme B
- IV. Biosynthesis of Methanofuran
- V. Biosynthesis of Coenzyme F₄₂₀
- VI. Biosynthesis of Coenzyme M
- References

Our current knowledge of the pathways and genes involved in the biosynthesis of the methanogenic coenzymes methanopterin, coenzyme B, methanofuran, coenzyme F₄₂₀, and coenzyme M is presented. Proposed reaction mechanisms for several of the novel reactions involved in the pathways are presented. © 2001 Academic Press.

I. INTRODUCTION

This review on the biosynthesis of the methanogenic cofactors in the methanoarchaea will constitute an update of published and unpublished work on this topic since it was last reviewed in 1993 (White and Zhou, 1993). The structures of the cofactors are shown in the figures and include methanopterin, coenzyme B, methanofuran, coenzyme F₄₂₀, and coenzyme M. These coenzymes are referred to as the methanogenic coenzymes, because each functions as a catalyst in the biochemical reduction of CO₂ or acetate to CH₄ (DiMarco *et al.*, 1990).

The present status of work on each of the coenzymes will be presented using the figures, which show the biosynthetic pathways, as the focus of the discussion. Most of the work reported involves the cloning and overexpression of *Methanococcus jannaschii* gene products. Because they are derived from the smallest hyperthermophilic autotrophic methanogenic archaeal genome currently sequenced, and therefore fewer genes have to be considered, identification of the desired genes is more likely. We have also confirmed that *M. jannaschii* contains the coenzymes that are discussed in this chapter (Graupner and White,

2000). Furthermore, purification of the overexpressed enzymes in *Escherichia coli* is easily accomplished since most of the *E. coli* proteins precipitate on heating, whereas the *M. jannaschii* proteins—being from a hyperthermophile—do not precipitate. Measurements of the biosynthetic reactions of the overexpressed enzymes have been performed with extracts of *Methanosarcina thermophila*, because these cells consistently show higher activities than those observed with other methanoarchaea.

II. BIOSYNTHESIS OF METHANOPTERIN

Methanopterin (MPT), in its tetrahydro reduced state, functions as a C_1 carrier coenzyme in methanogenesis (DiMarco *et al.*, 1990; Weiss and Thauer, 1993; Thauer, 1998) in a capacity analogous to that seen in folate biochemistry, where 5N -formyltetrahydrofolate is converted to 5N - ^{10}N -methylene-tetrahydrofolate and finally to 5N -methyltetrahydrofolate (Matthews, 1996). On the basis of chemical and stereochemical (White, 1996b) similarities between the structures of tetrahydrofolate and tetrahydromethanopterin, some common connections between the biosynthesis of these two cofactors were considered likely. These connections have been confirmed by a series of early labeling experiments, which show that GTP (I) is the precursor of the pterin and that 4-aminobenzoic acid is the precursor of the arylamine portion of the molecule (White and Zhou, 1993).

Of the seventeen reactions involved in the biosynthesis of the reduced form of methanopterin (XXI) (Fig. 1), all reactions but the one analogous to the dihydrofolate reductase, reaction 17, have been demonstrated to occur in cell extracts of *M. thermophila* (White, 1990, 1996a), an organism containing sarcinapterin, an analog of methanopterin. The steps in the biosynthesis of MPT in *Methanobacterium thermoautotrophicum* strains ΔH and Marburg have been shown to be essentially the same as those established for *M. thermophila*, demonstrating the universal occurrence of this pathway in the methanogens (White, 1998).

So far, only three genes in the Methanopterin biosynthetic pathway (Fig. 1) have been identified in *Methanococcus jannaschii*: MJ0798, MJ0301, and MJ1425. The enzyme derived from gene MJ0798 catalyzes reaction 1 in Fig. 1; that from MJ0301, reaction 7; and that from MJ1425, reaction 18. Each of these genes has been confirmed by the overexpression of the gene product and by establishing that it catalyzes the desired reaction.

H. WHITE

METHANOGENIC COFACTORS

301

f the overexpressed enzymes in *Es-*
d since most of the *E. coli* proteins
 he *M. jannaschii* proteins—being
 precipitate. Measurements of the
 xpressed enzymes have been per-
arcina thermophila, because these
 ities than those observed with oth-

OF METHANOPTERIN


hydro reduced state, functions as a
 nesis (DiMarco *et al.*, 1990; Weiss
 a capacity analogous to that seen
 myltetrahydrofolate is converted to
 and finally to ⁵N-methyltetrahy-
 basis of chemical and stereochem-
 between the structures of tetrahydrofo-
 some common connections between
 ctors were considered likely. These
 y a series of early labeling experi-
 the precursor of the pterin and that
 or of the arylamine portion of the

ed in the biosynthesis of the reduced
 1), all reactions but the one analo-
 ae, reaction 17, have been demon-
thermophila (White, 1990, 1996a),
 in, an analog of methanopterin. The
 in *Methanobacterium thermoau-*
rg have been shown to be essential-
M. thermophila, demonstrating the
 way in the methanogens (White,

ethanopterin biosynthetic pathway
ethanococcus jannaschii: MJ0798,
 ie derived from gene MJ0798 cat-
 MJ0301, reaction 7; and that from
 e genes has been confirmed by the
 and by establishing that it catalyzes

The biosynthetic pathway to MPT begins with the removal of the C-8 carbon of GTP (I) to form triaminopyrimidine triphosphate (PTP, II), a previously unidentified natural product. The removal of the C-8 carbon of GTP is a common feature of both the GTP cyclohydrolase I reaction, involved in neopterin biosynthesis (Green *et al.*, 1996), and the GTP cyclohydrolase II reaction, involved in riboflavin biosynthesis (Bacher *et al.*, 1996) in both bacteria and eukaryotic organisms. The reactions required for the archaeal transformation are consistent with the first two mechanistic steps involved in the bacterial GTP cyclohydrolase II reaction (Foor and Brown, 1975) but are not consistent with the GTP cyclohydrolase I mechanism (Nar *et al.*, 1995). The formation of compound II has been shown in our laboratory to be catalyzed by the enzyme encoded by the *M. jannaschii* gene MJ0798. Compound II is not an intermediate in the GTP cyclohydrolase I or II enzymatic reactions, but it is an intermediate in methanopterin biosynthesis in *M. thermophila* (Howell and White, 1997). Compound II would, however, appear to serve as a common precursor for methanopterin, riboflavin, and coenzyme F₄₂₀ in the methanoarchaea (see later). The methanoarchaeal GTP cyclohydrolase may in fact be related to GTP cyclohydrolase II, the first enzyme in bacterial riboflavin biosynthesis, since it has been identified by a 21-amino acid sequence with 25% identity to the *Escherichia coli* GTP cyclohydrolase II. The enzyme also has an 81-

FIG. 1. Pathway and genes involved in the biosynthesis of 5,6,7,8-tetrahydro-methanopterin (XXI). In all the figures included in this review, intermediates are indicated by roman numerals, reactions by underlined numbers, and reactions that have been confirmed in cell extracts and/or with the cloned gene products by circled underlined reaction numbers. Reactions for which the genes have been assigned are indicated by the *M. jannaschii* gene number or the gene name, if this has been assigned. The gene numbers refer to genes assigned in *M. jannaschii* (Bult *et al.*, 1997). I, Guanosine triphosphate; II, triaminopyrimidine triphosphate; III, 7,8-dihydroneopterin 2':3'-cyclic phosphate; IV, 7,8-dihydroneopterin 3'-phosphate; V, 7,8-dihydroneopterin; VI, glycolaldehyde; VII, 6-hydroxymethyl-7,8-dihydropterin; VIII, 6-hydroxymethyl-7,8-dihydropterin pyro-phosphate; IX, 4-(β-D-ribofuranosyl)aminobenzene 5'-phosphate; X, 5-phospho-β-D-ribose-1-pyrophosphate; XI, 4-aminobenzoic acid; XII, 7,8-dihydropterin-6-ylmethyl-4-(β-D-ribofuranosyl) aminobenzene 5'-phosphate; XIII, 7,8-dihydropterin-6-ylmethyl-1-(4-aminophenyl)-1-deoxy-D-ribitol 5'-phosphate; XIV, 7,8-dihydropterin-6-ylmethyl-1-(4-aminophenyl)-1-deoxy-D-ribitol; XV, 7,8-dihydropterin-6-ylmethyl-1-(4-aminophenyl)-1-deoxy-5-[1-α-D-ribofuranosyl 5'-diphosphate]-D-ribitol; XVI, 7,8-dihydropterin-6-ylmethyl-1-(4-aminophenyl)-1-deoxy-5-[1-α-D-ribofuranosyl 5'-triphosphate]-D-ribitol; XVII, 7,8-dihydropterin-6-ylmethyl-1-(4-aminophenyl)-1-deoxy-5-[1-α-D-ribofuranosyl 5'-triphosphate]-D-ribitol; XXII, α-ketoglutarate; XXIII, (S)-2-hydroxyglutaric acid; XVIII, didemethylated 7,8-dihydromethanopterin; XIX, monomethylated 7,8-dihydromethanopterin; XX, 7,8-dihydromethanopterin; XXI, 5,6,7,8-tetrahydromethanopterin.



amino acid sequence having 21% identity with the putatively identified *Archaeoglobus fulgidus* GTP cyclohydrolase II derived from AF0484 (Klenk *et al.*, 1998). We have named the *M. jannaschii* enzyme "GTP cyclohydrolase III" since the product of its enzymatic reaction, compound II, is different from that produced by either GTP cyclohydrolase I, which produces 7,8-dihydroneopterin triphosphate, or GTP cyclohydrolase II, which produces triaminopyrimidine monophosphate (I in Fig. 9) (Howell and White, 1997).

Compound II is then converted to 7,8-dihydroneopterin 2':3'-cyclic phosphate (III) (Fig. 1, reaction 2), an intermediate identified by incubating a cell extract of *M. thermophila* with GTP (Howell and White, 1997). A proposed mechanism for this reaction is shown in fig. 2. Hydrolysis of the cyclic phosphate in compound III (reaction 3 in Fig. 1) leads to 7,8-dihydroneopterin 3'-phosphate (IV), which is then hydrolyzed to 7,8-dihydroneopterin (V) by an unknown phosphatase (reaction 4 in Fig. 1).

The next two reactions in MPT biosynthesis (reactions 5 and 6 in Fig. 1) are identical to the reactions in folate biosynthesis and would be expected to be catalyzed by dihydroneopterin aldolase and by 6-hydroxymethyldihydropterin pyrophosphokinase, respectively. For both of these reactions, the bacterial genes are known (Lopez and Lacks, 1993; Talarico *et al.*, 1992), and the X-ray crystal structure for the enzyme catalyzing both reactions 5 and 6 has been reported (Hennig *et al.*, 1998, 1999; Xiao *et al.*, 1999; Stammers *et al.*, 1999). Despite this information, none of the genes for these enzymes can be found in the genomes of the archaea *M. jannaschii*, *M. thermoautotrophicum* strain ΔH , or *A. fulgidus*, based on sequence comparisons. It would appear that either nonorthologous enzymes are responsible for catalyzing these reactions in the methanoarchaea, or the sequences of the enzymes have diversified to such an extent that they cannot be detected by sequence comparisons.

The next reaction in the pathway (reaction 7 in Fig. 1) is the condensation of 6-hydroxymethyl-7,8-dihydropterin pyrophosphate (VIII) with 4-(β -D-ribofuranosyl)aminobenzene 5'-phosphate (β -RFA-P, IX) to form XII. This reaction is mechanistically analogous to the 7,8-dihydropteroate synthase in bacterial folate biosynthesis (Green *et al.*, 1996), but it utilizes IX in place of 4-aminobenzoic acid (XI) or *N*-(*p*-aminobenzoyl)-L-glutamic acid. Again, no gene with sequence homology to the 7,8-dihydropteroate synthase can be readily identified in the archaeal genomes. The gene encoding the enzyme has been identified however, using a novel computer program, ORF (an acronym for Ostensible Recognition of Folds), developed by Aurora and Rose (1998) for

L. WHITE

ntity with the putatively identified ydrolase II derived from AF0484 the *M. jannaschii* enzyme "GTP cyf its enzymatic reaction, compound I by either GTP cyclohydrolase I, in triphosphate, or GTP cyclohy-pyrimidine monophosphate (I in

o 7,8-dihydroneopterin 2':3'-cyclic an intermediate identified by incu-*tila* with GTP (Howell and White, is reaction is shown in fig. 2. Hy-ompound III (reaction 3 in Fig. 1) hosphate (IV), which is then hy-) by an unknown phosphatase (re-

synthesis (reactions 5 and 6 in Fig. late biosynthesis and would be ex-opterin aldolase and by 6-hydroxy-*cinase*, respectively. For both of are known (Lopez and Lacks, 1993; rystal structure for the enzyme cat-een reported (Hennig *et al.*, 1998, *al.*, 1999). Despite this information, is can be found in the genomes of *noautotrophicum* strain ΔH , or *A.* isons. It would appear that either sible for catalyzing these reactions ences of the enzymes have diversi-not be detected by sequence com-

(reaction 7 in Fig. 1) is the conden-ydropterin pyrophosphate (VIII) zene 5'-phosphate (β -RFA-P, IX) stically analogous to the 7,8-dihy-folate biosynthesis (Green *et al.*, 4-aminobenzoic acid (XI) or *N*-(*p*-in, no gene with sequence homolo-ase can be readily identified in the ng the enzyme has been identified rogram, ORF (an acronym for Os-oped by Aurora and Rose (1998) for





FIG. 2. Proposed intermediates in the conversion of triaminopyrimidine triphosphate (II) to 7,8-dihydroneopterin 2':3'-cyclic phosphate (III). II, Triaminopyrimidine triphosphate; III, 7,8-dihydroneopterin 2':3'-cyclic phosphate; XXV and XXVI are proposed intermediates in the pathway.

the identification of genes in genomic databases. This program can identify open reading frames with low sequence homologies based on their predicted secondary structure distributions when compared with secondary structure distributions of a known enzyme. The enzyme identified in this search, derived from *M. jannaschii* gene MJ0301, has only 19.9% sequence similarity to the fungal 7,8-dihydropteroate synthase from *Pneumocystis carinii*. The enzyme has been overexpressed and shown to catalyze reaction 7 (Fig. 1; Xu *et al.*, 1999). The enzyme does not catalyze the formation of 7,8-dihydropteroate from **XI** and **VIII**. This is also found to be true for gene product of MJ0107, which has been putatively assigned as a 7,8-dihydropteroate synthase (Bult *et al.*, 1997). The discovery that the *M. jannaschii* thymidylate synthase, derived from MJ0757, uses methylenetetrahydrofolate as the co-factor for the formation of dTMP (Xu *et al.*, 1999) is surprising because the methanobacteria are considered to lack folates (Leigh, 1983; Worrell and Nagle, 1988).

One of the unique steps in the MPT biosynthetic pathway (reaction 8 in Fig. 1) is the condensation of **XI** with 5-phospho- α -D-ribosyl diphosphate (PRPP, **X**) to produce **IX**. This reaction, catalyzed by the *M. thermophila* β -RFA-P synthase, is unique among known phosphoribosyltransferases (PRTases) in that a decarboxylation of one of the substrates (**XI**) occurs during the reaction, and a C-riboside rather than a N-riboside is formed at the site of the decarboxylation (Rasche and White, 1998). The enzymatic reaction is very specific for **XI**, with 4-hydroxybenzoic acid serving as the only alternate substrate. This reaction is, however, analogous to other PRTase reactions in that pyrophosphate is released as a product of the reaction and the reaction proceeds with a net inversion of stereochemistry at C-1 of the ribose, as shown in Fig. 3. Evidence for this mechanism was obtained by the NaCNBH₃ reduction of intermediate **I** to β -D-ribofuranosylbenzene 5'-phosphate (Rasche and White, 1998).

Compound **XII** is reduced to compound **XIII** (reaction 9 in Fig. 1) in a reaction stimulated by the addition of FMN or coenzyme F₄₂₀ to the reaction mixture (White, 1996a); as with reaction 10, no information is available concerning the enzymes or genes for these reactions. A possi-

FIG. 3. Mechanisms for reactions 8 and 11 (see Fig. 1) in methanopterin biosynthesis. **IX**, 4-(β -D-Ribofuranosyl)aminobenzene 5'-phosphate; **X**, 5-phospho- β -D-ribosyl-1-pyrophosphate; **XI**, 4-aminobenzoic acid; **XXVII**, oxonium ion formed by the loss of pyrophosphate from PRPP; **XIV**, 7,8-dihydropterin-6-ylmethyl-1-(4-aminophenyl)-1-deoxy-D-ribose; **XV**, 7,8-dihydropterin-6-ylmethyl-1-(4-aminophenyl)-1-deoxy-5-[1- α -D-ribofuranosyl 5'-phosphate]-D-ribose; **XXVIII**, intermediate **I**.



ble mechanism for this reaction would involve the intramolecular elimination of an alcohol in **XII** to form a substituted methylene quinone imine (**XXIX**) that would then be reduced to compound **XIII** (Fig. 4).

Reaction 11 in Fig. 3 is another unique biochemical reaction for a PRTase since the product of the reaction, **XV** (**XIII** with an α -linked ribose phosphate in place of the terminal phosphate), contains an α -linked ribosyl group. All previous known examples of PRTase products





FIG. 4. Mechanisms for reactions 9, 12, 13, and 14 (see Fig. 1) in methanopterin biosynthesis. **XII**, 7,8-Dihydropterin-6-ylmethyl-4-(β -D-ribofuranosyl)aminobenzene 5'-phosphate; **XIII**, 7,8-dihydropterin-6-ylmethyl-1-(4-aminophenyl)-1-deoxy-D-ribitol 5'-phosphate; **XV**, 7,8-dihydropterin-6-ylmethyl-1-(4-aminophenyl)-1-deoxy-5-[1- α -D-ribofuranosyl 5'-phosphate]-D-ribitol; **XVI**, 7,8-dihydropterin-6-ylmethyl-1-(4-aminophenyl)-1-deoxy-5-[1- α -D-ribofuranosyl 5'-diphosphate]-D-ribitol; **XVII**, 7,8-dihydropterin-6-ylmethyl-1-(4-aminophenyl)-1-deoxy-5-[1- α -D-ribofuranosyl 5'-triphosphate]-D-ribitol; **XXIII**, (S)-2-hydroxyglutaric acid; **XVIII**, didemethylated 7,8-dihydropterin.

unique biochemical reaction for a reaction, XV (XIII with an α -linked triminal phosphate), contains an α -known examples of PRTase products



13, and 14 (see Fig. 1) in methanopterin methyl-4-(β -D-ribofuranosyl)aminobenzene methyl-1-(4-aminophenyl)-1-deoxy-D-ribitol ethyl-1-(4-aminophenyl)-1-deoxy-5-[1- α -D-7,8-dihydropterin-6-ylmethyl-1-(4-aminophosphate)-D-ribitol; XVII, 7,8-dihydropterin-5-[1- α -D-ribofuranosyl 5'-triphosphate)-D-ribose, didemethylated 7,8-dihydromethanopterin.

Labeling experiments with cell extracts of *M. thermoautotrophicum* strain ΔH and Marburg and *M. thermophila* strain TM-1 have established that the sites in the overall sequence of reactions from II to 5,6,7,8-tetrahydromethanopterin (XXI)—where the *S*-adenosylmethionine (SAM)-dependent C-9 and C-7 methylations of the pterin-containing intermediates occur—are dependent on the species of methanoarchaea (White, 1998). I have proposed that MJ0563 and/or MJ1200, putatively identified as DNA-cytosine methyltransferases, may be the genes encoding the enzymes responsible for these methylation reactions (reactions 15 and 16 in Fig. 1) (White, 1998). This idea is supported by the observation that these methanoarchaea do not contain 5-methylcytosine-modified bases (J. A. McClosky, personal communication, 1996). Since the methylations are sensitive to thiol reagents, a mechanism requiring the addition of the thiol to the different enamine isomers of the dihydropterin, as shown in Fig. 5, has been proposed (White, 1998). This reaction mechanism is analogous to that proposed for thymidylate synthase (Carreras and Santi, 1995) and 2'-deoxyuridylate hydroxymethylase (Kunitani and Santi, 1980). It has also been demonstrated during this work that cells of *M. thermoautotrophicum* strains ΔH and Marburg contain significant amounts of methanopterin that lacks the phosphate and 2-hydroxyglutaric acid groups (White, 1998).

Given that the absolute stereochemistry of the hydroxyglutaric acid




FIG. 6. Proposed mechanism for the methylation of the intermediates leading to methanopterin. The chemical structure(s) of the specific methanopterin intermediates that are methylated is species dependent. Thus the R group can represent several different structures.




FIG. 6. Proposed mechanism for the methylation of the intermediates leading to methanopterin. The chemical structure(s) of the specific methanopterin intermediates that are methylated is species dependent. Thus the R group can represent several different structures.

present in MPT has been established as (S)-hydroxyglutaric acid (XXI-II) (Solow and White, 1997a), it is considered likely that this moiety of MPT is produced by the reduction of α -ketoglutaric acid by a lactate/malate dehydrogenase-like enzyme. The *M. jannaschii* genome contains two genes predicted to produce homologs of the lactate/malate dehydrogenase group of enzymes, MJ0490 and MJ1425. We have cloned and overexpressed the enzyme derived from each of these genes and have determined that the protein derived from MJ1425 functions as an α -ketoglutarate reductase, producing XXIII by the NADH-dependent reduction of α -ketoglutarate (XXII) (Graupner *et al.*, 2000). Thus, this enzyme is very likely involved in the production of the XXIII moiety of the side chain of methanopterin (reaction 18 in Fig. 1). This same enzyme also reduces sulfolpyruvate to L-sulfolactate and thus functions in the biosynthesis of coenzyme M (see later).

Many additional analogs of methanopterin function throughout the archaea (White, 1993a,b, 1997), and methanopterin has now been found in organisms outside of the archaeal domain (Chistoserdova *et al.*, 1998; Pomper *et al.*, 1999). Since the structures of all of these modified folates have common features, they will surely be found to have related biosynthetic pathways.

III. BIOSYNTHESIS OF COENZYME B

The biochemistry of the steps involved in the conversion of α -ketoglutarate and acetyl coenzyme A (acetyl-CoA) to α -ketosuberate, a precursor of coenzyme B [7-mercaptoheptanoylthreonine phosphate (CVII)] and of biotin, in *Methanosarcina thermophila*, and its further conversion to coenzyme B, is shown in Fig. 6. Most of the enzymes or genes involved in the pathway are now established and are indicated in Fig. 6. The biosynthesis of the alkyl portion of coenzyme B and biotin in the archaea is an interesting dilemma, considering that these cells contain no fatty acids (Kates, 1993) and have no genes homologous to the fatty acid synthases (Bult *et al.*, 1997).

Several years ago, a pathway for the formation of the 7-mercaptoheptanoic acid (CV) moiety of coenzyme B (CVII) was proposed based on ^{13}C -labeling studies and the identification of α -ketoglutarate (XXII), α -ketoadipate (XCV), α -ketopimelate (XCIX), and α -ketosuberate (CIII) in a series of methanoarchaea (White, 1989a,b). In the proposed pathway the repeated application of the α -ketoacid chain elongation series of reactions was used to increase the number of methylenes from two, as found in XXII, to five, as found in the product, CIII. Compound CIII





Fig. 6. Pathway and genes involved in the biosynthesis of coenzyme B and biotin. **XXII**, α -Ketoglutarate, 2-oxopentanedioic acid; **XC**, acetyl coenzyme A; **XCI**, *trans*-homoaconitate, (E)-1,2,4-but-1-enetricarboxylic acid; **XCIH**, (S)-homocitrate; (S)-2-hydroxy-1,2,4-butanetricarboxylic acid; **XCHH**, *cis*-homoaconitate, (Z)-1,2,4-but-1-enetricarboxylic acid; **XCIIV**, (-)-*threo*-isohomocitrate, (2R,3S)-1-hydroxy-1,2,4-butanetricarboxylic acid; **XCV**, α -ketoadipate, 2-oxohexanedioic acid; **XCVI**, (R)-(*homo*)-citrate, (R)-2-hydroxy-1,2,5-pentanetricarboxylic acid; **XCVII**, *cis*-(*homo*)-aconitate, (Z)-1,2,5-pent-1-enetricarboxylic acid; **XCVIII**, (-)-*threo*-iso(*homo*)-citrate, (2R,3S)-1-hydroxy-1,2,5-pentanetricarboxylic acid.



Knowing that the precursor to the α -ketoadipate (XCV) is XCVI, the same isohomocitrate isomer that is involved in lysine biosynthesis, and knowing that only trans additions and eliminations of water are involved in the interconversions of the intermediates (Gawron and Ma-

acid; **XCIX**, α -ketopimelate, 2-oxoheptanedioic acid; **C**, (*R*)-(homo)₅citrate, 2-hydroxy-1,2,6-hexanetricarboxylic acid; **CL**, *cis*-(homo)₄aconitate, (2*Z*)-1,2,6-hex-1-enetricarboxylic acid; **CH**, (–)-*threo*-isat(homo)₄citrate, (2*R*,3*S*)-1-hydroxy-1,2,6-hexanetricarboxylic acid; **CIH**, α -ketosuberate, 2-oxooctanedioic acid; **CIV**, 7-oxoheptanoic acid; **CV**, 7-mercaptoheptanoic acid; **CVI**, 7-mercaptoheptanoylthreonine; **CVII**, coenzyme B, 7-mercaptoheptanoylthreonine phosphate.

hajan, 1966), the reaction sequences 2, 3, and 4 shown in Fig. 6 have been proposed to account for the first set of reactions in the pathway. This pathway accounts for the formation of all of the intermediates observed. Reactions 2 and 3 would not be expected to be carried out by an enzyme homologous to aconitase or homoaconitase, since these enzymes use only **XCIII** and (*R*)-homocitrate as substrates (Beinert *et al.*, 1996). This idea has been strengthened by demonstrating that the two subunit aconitases formed by the different combinations of the proteins encoded by *M. jannaschii* genes MJ1003, MJ1271, MJ0499, and MJ1277 have not been found to catalyze reactions 2 and 3. Reaction 4, however, is that expected for an aconitase or a homoaconitase-like enzyme and is catalyzed by mixtures of these enzymes. It is possible that reaction 3 in Fig. 6 is catalyzed by the aconitate hydratase described by Neilson (1956a,b). Unfortunately, this enzyme appears to have never been purified nor isolated in a pure form; therefore, no information is currently available as to the stereospecificity of its reaction mechanism. Reaction 2 in Fig. 6 could be catalyzed by an enzyme related to mesaconase, an enzyme that catalyzes the reversible addition of water to mesaconate (*trans*-1,2-prop-1-enedicarboxylic acid) to form (*S*)-citramalate (Flint and Allen, 1996; Wang and Barker, 1969). The direct conversion of **XCI** to **XCIII** with an enzyme analogous to aconitate isomerase is not possible on mechanistic grounds (Klinman and Rose, 1971).

Rehydration of **XCIII** produces (*-*)-*threo*-isohomocitrate (**XCIV**), which undergoes a NAD-dependent oxidative decarboxylation to produce **XCV**. This reaction, which is analogous to an isocitrate dehydrogenase reaction, has been shown to be catalyzed by the protein product of the *M. jannaschii* gene MJ1596 (Howell *et al.*, 2000b). This gene product, now designated as AksF, and its homolog MJ0940—both originally assigned as isocitrate dehydrogenase (Bult *et al.*, 1997)—in fact do not catalyze the isocitrate dehydrogenase reaction with isocitrate as a substrate. The **XCV** resulting from this reaction then undergoes two consecutive sets of α -ketoacid chain elongation reactions to produce **CIII**. In each of these sets of reactions, it has been shown that the homologs of *cis*-homoaconitate, homocitrate, and (*-*)-*threo*-isohomocitrate serve as intermediates. The gene product of MJ0503 also catalyzes the condensation of **XCV** or **XCIX** with acetyl-CoA to form the (*R*)-homocitrate homologs of (*R*)-2-hydroxy-1,2,5-pentanetricarboxylic acid and of (*R*)-2-hydroxy-1,2,6-hexanetricarboxylic acid, respectively. The aconitase-like enzyme formed from the protein product of MJ1003 and MJ1271, designated AksDE, catalyzes reactions 4, 7, 8, 11, and 12 (see Fig. 6).

Compound **CIII** resulting from this series of reactions then undergoes a nonoxidative decarboxylation to form 7-oxoheptanoic acid (**CIV**)

s 2, 3, and 4 shown in Fig. 6 have set of reactions in the pathway. tion of all of the intermediates ob- be expected to be carried out by an r homoaconitase, since these en- itrate as substrates (Beinert *et al.*, ed by demonstrating that the two erent combinations of the proteins 003, MJ1271, MJ0499, and MJ1277 ions 2 and 3. Reaction 4, however, homoaconitase-like enzyme and is ies. It is possible that reaction 3 in te hydrase described by Neilson ie appears to have never been pu- erefore, no information is current- y of its reaction mechanism. Reac- an enzyme related to mesaconase, le addition of water to mesaconate l) to form (S)-citramalate (Flint and). The direct conversion of XCI to aconitate isomerase is not possible d Rose, 1971).

(-)-threo-isohomocitrate (XCIV), oxidative decarboxylation to pro- alogous to an isocitrate dehydro- e catalyzed by the protein product well *et al.*, 2000b). This gene prod- homolog MJ0940—both original- ase (Bult *et al.*, 1997)—in fact do nase reaction with isocitrate as a this reaction then undergoes two ngation reactions to produce CIII. is been shown that the homologs of (-)-threo-isohomocitrate serve as J0503 also catalyzes the condensa- A to form the (R)-homocitrate ho- etricarboxylic acid and of (R)-2-hy- respectively. The aconitase-like uct of MJ1003 and MJ1271, desig- , 8, 11, and 12 (see Fig. 6).

is series of reactions then under- to form 7-oxoheptanoic acid (CIV)

(reaction 15), a precursor to coenzyme B, and an oxidative decarboxy- lation to form pimeloyl-CoA (reaction 14), the precursor to biotin. The 7-oxo group of the 7-oxoheptanoic acid is then converted to a thiol by a previously described route (reaction 16) (White, 1989c). The details of the mechanism of this reaction are not presently understood, but they involve a protein-bound sulfur, derived from H₂S, as the source of the sulfur. The resulting 7-mercaptoheptanoic acid (CV) is converted, by an ATP-dependent process, to 7-mercaptoheptanoyl-L-threonine (CVI) (re- action 17) (Solow and White, 1997b). The current evidence indicates that this reaction proceeds by the formation of 7-mercaptoheptanoyl phosphate by the transfer of the γ -phosphate of the ATP to CV. Reac- tion of the amino group of L-threonine with the 7-mercaptoheptanoyl phosphate produces CVI. ATP-dependent phosphorylation of the CVI produces the final coenzyme B (reaction 18 in Fig. 6) (White, 1994). Growing cells of both *Methanococcus volta* and *M. thermophila* have been found to readily incorporate CV and CVI into coenzyme B. In the case of CVI the incorporation proceeds without cleavage of the amide bond (White, 1994).

IV. BIOSYNTHESIS OF METHANOFURAN

The most recent investigation relating to the biosynthesis of methanofu- ran has concentrated on the biosynthesis of the 1,3,4,6-hexanete- tracarboxylic acid (HTCA) portion of methanofuran (LXXXIII). A sum- mary of the present status of this work is shown in Fig. 7 and is in agreement with earlier labeling studies (Eisenreich and Bacher, 1992; Gorkovenko *et al.*, 1994; White, 1987). All of the circled reactions shown in the pathway have been confirmed by the synthesis of each interme- diate and by the demonstration that each is converted to one or more of the indicated intermediates, using cell-free preparations of *M. ther- mophila* (Howell *et al.*, 2000a). The pathway proposed for the biosyn- thesis of HTCA emerges in a logical manner from the intermediates de- tected in the methanogens, from the intermediates shown to be metabolized by cell extracts of *M. thermophila*, and from known infor- mation about biochemical reactions. The unusual aspect of the pathway is the cooccurrence of two different stereoisomers of the same compound functioning in a parallel manner in many of the steps in the pathway.

The first step in the pathway is the synthesis of *trans*-homoaconitate (XCI) from XXII and XC under the direction of the *aksA* gene product AksA (Fig. 7, reaction 1). This step is also the first step in the biosyn- thesis of α -ketosuberate, a precursor to the 7-mercaptoheptanoyl moi-

ety of coenzyme B, as discussed earlier in this chapter. Compound **XCI** thus represents a branch point metabolite leading to both coenzyme B and methanofuran. Trans addition of water to **XCI** produces an *erythro*-isohomocitrate (**LX**) (Fig. 7, reaction 2). This reaction involves the same trans addition that occurs with aconitase and fumarase enzymes, but it proceeds with a trans substrate analogous to that observed with the enzyme mesaconase (Flint and Allen, 1996; Wang and Barker, 1969). Although there is currently no protein sequence information available for any mesaconase, the genomes of *M. jannaschii* (Bult *et al.*, 1997), *M. thermoautotrophicum* ΔH (Smith *et al.*, 1997), and *A. fulgidus* (Klenk *et al.*, 1998) each have two sets of genes (Discussed in Section III) homologous to an aconitase or isopropylmalate dehydratase, with one member of each set containing the required [4Fe-4S] binding sites. We have cloned and overexpressed the enzymes from each of these *M. jannaschii* genes and have demonstrated that no combination of any of these proteins catalyzes any of the dehydration reactions involved in HTCA biosynthesis. The enzyme catalyzing this reaction is, however, not unique to the archaea since *E. coli* cell extracts have also been shown to readily catalyze this reaction (R. H. White, unpublished results, 1998). At present it has not been possible to establish which of the enantiomers of *erythro*-isohomocitrate is used in this pathway, since no method has been devised for the synthesis or isolation of the different enantiomers of **LX**. However, this is the first report of the occurrence and metabolism of an *erythro* isomer of isohomocitrate in nature.

The conversion of **LX** to the two different diastereomers of 1-hydroxy-1,3,4,5-pentanetetracarboxylic acid, **LXIII** and **LXIV** (Fig. 7, reaction 4), can be viewed as taking place in two separate steps. The first step consists of the oxidation of the hydroxyl group of **LX** to a keto group to produce oxaloglutarate, which then condenses with **XC** to form **LXIII** and **LXIV** in the second step. These reactions would be analogous to those used by the combined action of malate dehydrogenase (EC 1.1.1.37) and citrate synthase (EC 4.1.3.7). So far, it has not been possible to show that oxaloglutarate is an intermediate in this reaction. We therefore propose that the oxaloglutarate is a protein-bound intermediate as occurs in the isocitrate dehydrogenase-catalyzed reactions (Siebert *et al.*, 1957). An unusual aspect of the *erythro*-isohomocitrate reaction is that at least two different isomers are produced in the incubation, **LXIII** and **LXIV**. The production of these two isomers could be the result of two separate enzymes, one using (*S*)-oxaloglutarate (**LXI**) and the other using (*R*)-oxaloglutarate (**LXII**), or, more likely, it results by a single enzyme that is not sensitive to the stereochemistry at C-3 of **LXI** or **LXII** when carrying out reaction 4. It is interesting to note

WHITE

r in this chapter. Compound XCI olite leading to both coenzyme B vater to XCI produces an *erythro*-1. This reaction involves the same tase and fumarase enzymes, but dologous to that observed with the 1996; Wang and Barker, 1969). 1 sequence information available *jannaschii* (Bult *et al.*, 1997), *M. l.*, 1997), and *A. fulgidus* (Klenk ies (Discussed in Section III) ho- ymalate dehydratase, with one uired [4Fe-4S] binding sites. We zymes from each of these *M. jan*- d that no combination of any of ehydration reactions involved in lyzing this reaction is, however, oli cell extracts have also been n (R. H. White, unpublished re- m possible to establish which of ate is used in this pathway, since nthesis or isolation of the differ- s is the first report of the occur- mer of isohomocitrate in nature. rent diastereomers of 1-hydroxy- XIII and LXIV (Fig. 7, reaction wo separate steps. The first step yl group of LX to a keto group to ndenses with XC to form LXIII reactions would be analogous to of malate dehydrogenase (EC .3.7). So far, it has not been pos- intermediate in this reaction. We rate is a protein-bound interme- ydrogenase-catalyzed reactions ect of the *erythro*-isohomocitrate somers are produced in the incu- on of these two isomers could be e using (*S*)-oxaloglutarate (LXI) (LXII), or, more likely, it results ve to the stereochemistry at C-3 iction 4. It is interesting to note


METHANOGENIC COFACTORS

317

that the C-3 hydrogen of oxaloglutarate is expected to be acidic, lead- ing to rapid racemization at this carbon. This fact could account for the mixture of reaction products observed. The production of LXIII and LXIV now generates a branch point in the biosynthetic pathway where these two different isomers are metabolized to the final HTCA product.

The conversion of LXIII to (*E*)-1,3,4,5-pent-2-enetetra-carboxylic acid (LXV) and that of LXIV to (*Z*)-1,3,4,5-pent-2-enetetra-carboxylic acid (LXVII) are simple dehydration reactions. Each reaction could be cat- alyzed by an enzyme with a mechanism analogous to that observed in either an aconitase- or a fumarase-type reaction and would be expect- ed to proceed by the trans elimination of water. The production of (*Z*)- 1,3,4,5-pent-1-enetetra-carboxylic acid (LXVI) is also observed in these experiments and could arise by the trans elimination of water from the 1,2 positions of either LXIII or LXIV. Although most fumarases are rel- atively specific for fumarate, hydration of a series of substituted fu- marates has been observed by Eck and Simon (1994b), and a related en- zyme could be involved in the proposed pathway. The dehydration reaction could have been carried out by one of the aconitases present in *M. jannaschii* discussed earlier, but this has not been found to be the case.

FIG. 7. Pathway for the biosynthesis of the 1,3,4,6-hexanetetra-carboxylic acid moiety of methanofuran. XXII, α -Ketoglutarate; XCI, *trans*-homoaconitate, (*E*)-1,2,4-but-1-en- etricarboxylic acid; LX, *erythro*-isohomocitrate, (1*SR*,2*SR*)-1-hydroxy-1,2,4-butanetri- carboxylic acid; LXI, (*S*)-oxaloglutarate, (*S*)-1-oxo-1,2,4-butanetricarboxylic acid; LXII, (*R*)-oxaloglutarate, (*R*)-1-oxo-1,2,4-butanetricarboxylic acid; LXIII, *erythro*-HPTCA, (2*SR*,3*SR*)-2-hydroxy-1,2,3,5-pentanetetra-carboxylic acid; LXIV, *threo*-HPTCA, (2*SR*, 3*RS*)-2-hydroxy-1,2,3,5-pentanetetra-carboxylic acid; LXV, (*E*)-2-PETCA, (*E*)-1,2,3,5- pent-2-enetetra-carboxylic acid; LXVI, (*Z*)-1-PETCA, (*Z*)-1,2,3,5-pent-1-enetetra-car- boxylic acid; LXVII, (*Z*)-2-PETCA, (*Z*)-1,2,3,5-pent-2-enetetra-carboxylic acid; LXVIII, *L*-*erythro*-PTCA, *L*-*erythro*-(2*R*,3*S*)-1,2,3,5-pentanetetra-carboxylic acid; LXIX, *L*-*threo*- PTCA, *L*-*threo*-(2*S*,3*S*)-1,2,3,5-pentanetetra-carboxylic acid; LXX, coenzyme A ester of *L*-*erythro*-(2*R*,3*S*)-1,2,3,5-pentanetetra-carboxylic acid; LXXI, *L*-*xylo*-HHTCA, (1*S*,3*S*,4*S*)- 1-hydroxy-1,3,4,6-hexanetetra-carboxylic acid; LXXII, coenzyme A ester of *L*-*threo*- (2*S*,3*S*)-1,2,3,5-pentanetetra-carboxylic acid; LXXIII, *L*-*erythro*-KHTCA, (3*R*,4*S*)-1-oxo-1, 3,4,6-hexanetetra-carboxylic acid; LXXIV, *L*-*threo*-KHTCA, (3*S*,4*S*)-1-oxo-1,3,4,6-hexa- netetra-carboxylic acid; LXXV, *L*-*arabino*-HHTCA, (1*R*,3*R*,4*S*)-1-hydroxy-1,3,4,6-hexa- netetra-carboxylic acid; LXXVI, *L*-*ribo*-HHTCA, (1*S*,3*R*,4*S*)-1-hydroxy-1,3,4,6-hexanete- tracarboxylic acid; LXXVII, *L*-*xylo*-HHTCA, (1*R*,3*S*,4*S*)-1-hydroxy-1,3,4,6-hexanete- tracarboxylic acid; LXXVIII, coenzyme A ester of (1*R*,3*R*,4*S*)-1-hydroxy-1,3,4,6-hexanete- tracarboxylic acid; LXXIX, coenzyme A ester of (1*S*,3*R*,4*S*)-1-hydroxy-1,3,4,6-hexanete- tracarboxylic acid; LXXX, *L*-*erythro*-(*E*)-1,3,4,6-hex-1-enetetra-carboxylic acid, (3*R*,4*S*)- HETCA; LXXXI, *L*-*threo*-(*E*)-1,3,4,6-hex-1-enetetra-carboxylic acid, (3*S*,4*S*)-HETCA; LXXXII, coenzyme A ester of HTCA, (2*RS*,3*SR*)-1,3,4,6-hexanetetra-carboxylic acid or *meso*-HTCA; LXXXIII, methanofuran.







Fig. 7. See p. 317 for legend.

The hydrogenation of **LXV** to *L-erythro*-1,3,4,5-pentanetetracarboxylic acid (**LXVIII**) and of **LXVII** to *L-threo*-1,3,4,5-pentanetetracarboxylic acid (**LXIX**) (Fig. 7, reactions 7 and 8) would both be expected to proceed by a trans hydrogenation, as observed in the reduction of fumarate (Tchen and van Milligan, 1960), substituted fumarates (Eck and Simon, 1994a), and all known examples of the biochemical reduction of monounsaturated acids (White, 1980). The archaeal reductions (reactions 7 and 8) are most analogous to those carried out by either fumarate reductase or succinate dehydrogenase, which are closely related enzymes (van Hellemond and Tielens, 1994). The thiol:fumarate reductase for *M. thermoautotrophicum* ΔH (MT1850) (Heim *et al.*, 1998) has not been found to catalyze the reduction of a mixture of the 1,3,4,5-pentanetetracarboxylic acid isomers (R. Hedderich, personal communication, 1999). We propose a single enzyme for steps 7 and 8, which would catalyze the addition of hydrogen to the C-3 *re* face of either **LXV** or **LXVII** substrates. In this manner, the absolute stereochemistry of this carbon would be fixed in the required (*S*) configuration for all subsequent intermediates in this pathway (see later discussion).

We propose that **LXVIII** and **LXIX** are each converted to their respective CoA thioesters, **LXX** and **LXXII** (Fig. 7, reactions 9 and 10), followed by reductive carboxylation to 1-keto-1,3,4,6-hexanetetracarboxylic acid (KHTCA) isomers **LXXIII** and **LXXIV** (Fig. 7, reactions 11 and 12). Reaction 9 has been confirmed by the demonstration that **LXVIII** is converted to both **LXXIII** and **LXXIV** when incubated with HSCoA, HCO_3^- , and ATP. The detection of both **LXXIII** and **LXXIV**, which would result from epimerization of one or more of the asymmetric centers of the KHTCA molecule, was unexpected. The reason for this epimerization has been traced to the rapid exchange of the C-3 hydrogens of KHTCA with the protons of water. This exchange of the C-3 hydrogens of KHTCA is explained by the formation of the enol of the keto acid, which generates a vinylogous analog of malonic acid and as a result would be expected to have acidic methylene hydrogens leading to epimerization at C-3.

The formation of the CoA thioester of **LXVII** and **LXIX** required for steps 9 and 10 (and possibly also for steps 15 and 16 discussed later) could be carried out by several different routes that have been identified in the archaea. These routes could involve the sequential action of enzymes related to acetate kinase and phosphotransacetylase (Latimer and Ferry, 1993), an enzyme related to acetyl-CoA synthase (ADP-forming) (EC 6.2.1.1.3) (Glasemacher *et al.*, 1997), an enzyme related to the coenzyme A-transferases (Jacob *et al.*, 1997), or more likely an enzyme related to acetyl-CoA synthase (Eggen *et al.*, 1991; Jetten *et al.*, 1989).

-erythro-1,3,4,5-pentanetetra-carboxylic acid (L-threo-1,3,4,5-pentanetetra-carboxylic acid) 7 and 8) would both be expected as observed in the reduction of fumarate, substituted fumarates (Eck and Jørgensen, 1994). The archaeal reductions (reactions 9 and 10) are those carried out by either fumarate reductase, which are closely related to the bacterial ones (Jørgensen, 1994). The thiol:fumarate reductase (MT1850) (Heim *et al.*, 1998) reduces a mixture of the 1,3,4,5-tetrahydroxy-2-ketopentanoate (LXXIII) and LXXIV, which are the C-3 *re* face of either LXV or LXVI, the absolute stereochemistry of which is (S) configuration for all substrates (see later discussion).

X are each converted to their respective ketone (Fig. 7, reactions 9 and 10), to 1-keto-1,3,4,6-hexanetetra-carboxylic acid and LXXIV (Fig. 7, reactions 11 and 12). The demonstration that LXXIII and LXXIV when incubated with fumarate, both of which are the C-3 *re* face of either LXV or LXVI, on of one or more of the asymmetric carbons is unexpected. The reason for this rapid exchange of the C-3 hydroxyl group is not clear. This exchange of the C-3 hydroxyl group is the formation of the enol of the keto analog of malonic acid and as a result methylene hydrogens leading to

of LXVII and LXIX required for steps 15 and 16 discussed later) and the two different routes that have been identified involve the sequential action of 1) phosphotransacetylase (Latimer and Woese, 1997), an enzyme related to the bacterial one (Jørgensen, 1997), or more likely an enzyme (Jørgensen *et al.*, 1991; Jetten *et al.*, 1989).

The exact route used for the formation of the coenzyme A thioesters in KHTCA biosynthesis in the methanoarchaea has yet to be established.

The reductive carboxylation of acyl-CoA thioesters to α -ketoacids is a well-characterized reaction in the archaea. This reaction is known to be catalyzed by α -ketoacid ferredoxin oxidoreductases (α -ketoacid oxidoreductases) (Bock *et al.*, 1996; Heider *et al.*, 1996; Kletzin and Adams, 1996; Tersteegen *et al.*, 1997; Yoon *et al.*, 1996), which are not restricted to the archaea (Gehring and Arnon, 1972). These enzymes catalyze the reversible conversion of acyl-CoA thioesters and CO_2 to α -ketoacids and HSCoA in the presence of low-potential electron donors/acceptors. These enzymes contain thiamine pyrophosphate (TPP) and Fe/S centers and appear to proceed by a radical mechanism, as shown in Fig. 8 (Menon and Ragsdale, 1997). The largest group of α -ketoacid oxidoreductases have a tetrameric $\alpha\beta\gamma\delta$ structure, but through gene fusions of the subunit genes, enzymes with only one, two, and three subunits have also been identified (Bock *et al.*, 1996; Kletzin and Adams, 1996; Tersteegen *et al.*, 1997). Several sets of genes for this type of enzyme are present in *M. jannaschii*, *M. thermoautotrophicum* Δ H, and *A. fulgidus* genomes, one set of which is likely to be involved in catalyzing the required carboxylation of LXVIII and LXIX in the biosynthesis of KHTCA.

Incubation of a cell-free extract of *M. thermophilus* with a mixture of the KHTCA isomers, NADH, and NADPH produces about equal amounts of the *arabino*- and *xylo*-1-hydroxy-1,3,4,6-hexanetetra-carboxylic acid (HHTCA) diastereomers. This reduction of the KHTCA isomers (Fig. 7, reactions 13 and 14) could proceed either by a NADH- or a NADPH-dependent reduction, such as occurs with the pyridine nucleotide-dependent 2-hydroxyacid dehydrogenases such as lactate or malate dehydrogenase (Kim and Whitesides, 1988), or by a non-pyridine nucleotide-dependent 2-ketoacid reductase (Neumann and Simon, 1984; Simon *et al.*, 1985). Currently nothing is known about the genes for the non-pyridine nucleotide-dependent 2-ketoacid reductase. Two malate dehydrogenases (Mdh), designated Mdh I and Mdh II, however, have been isolated from *M. thermoautotrophicum* strain Marburg and, based on their N-terminal sequences, have most likely been encoded from the *M. thermoautotrophicum* strain Δ H genes MT1205 and MT0188, respectively (Thompson *et al.*, 1998).

We have cloned and overexpressed the proteins derived from the *M. jannaschii* genes MJ1425 and MJ0490 and have demonstrated that they catalyze the NADH/NADPH reduction of a series of different α -ketoacids to (S)-2-hydroxyacids. The enzyme from MJ1425, Mdh I, has been found to reduce the equilibrium mixture of the KHTCA isomers to only the *xylo* isomer of HHTCA (LXXI). On the basis of the stereospecificity of

322

ROBERT H. WHITE





FIG. 8. Reaction mechanisms for reactions 11 and 12 (top) and 17 and 18 (bottom) in the biosynthesis of the 1,3,4,6-hexanetetra-carboxylic acid moiety in methanofuran (see Fig. 7). TPP, Thiamine pyrophosphate; LXX, coenzyme A ester of *L-erythro*-(2*R*,3*S*)-1,2,3,5-pentanetetra-carboxylic acid; LXXII, coenzyme A ester of *L-threo*-(2*S*,3*S*)-1,2,3,5-pentanetetra-carboxylic acid; LXXXVIII, coenzyme A ester of (1*R*,3*R*,4*S*)-1-hydroxy-1,3,4,6-hexanetetra-carboxylic acid; LXXXIX, coenzyme A ester of (1*S*,3*R*,4*S*)-1-hydroxy-1,3,4,6-hexanetetra-carboxylic acid; LXXXV, anion of hemithioacetal intermediate; LXXXVI, hydroxyethyl-TPP radical intermediate; LXXXVII, carbanion of hydroxyethyl-TPP intermediate; LXXXVIII, hydroxyethyl-TPP intermediate; LXXXIX and XCI, ketyl radical intermediates; XC, enoxy radical intermediate.

this enzyme, this isomer is assigned as *L-xylo*-HHTCA (LXXI) (see Fig. 7 for the absolute stereochemistry and structure of these compounds). Since this isomer has not been found to be an intermediate in HTCA biosynthesis, whereas *D,L-arabino*-HHTCA has been found to be an intermediate, it is concluded that *L-arabino*-HHTCA (LXXV) and *L-lyxo*-HHTCA (LXXXVII) are the true intermediates in the pathway. Since

WHITE

METHANOGENIC COFACTORS

323



11 and 12 (top) and 17 and 18 (bottom) in coxylic acid moiety in methanofuran (see , coenzyme A ester of *L-erythro*-(2*R*,3*S*)-coenzyme A ester of *L-threo*-(2*S*,3*S*)-1,2,3,5-oxime A ester of (1*R*,3*R*,4*S*)-1-hydroxy-coenzyme A ester of (1*S*,3*R*,4*S*)-1-hydroxy-1,1 of hemithioacetal intermediate; LXXXVI, XVII, carbanion of hydroxyethyl-TPP intermediate; LXXXIX and XCI, ketyl radi-

is *L-xylo*-HHTCA (LXXI) (see Fig. id structure of these compounds). I to be an intermediate in HTCA HTCA has been found to be an *in-bino*-HHTCA (LXXV) and *L-lyxo*-rmediates in the pathway. Since

there is no mechanism for the stereochemistry at C-3 to change in any intermediate in the pathway after LXXV, LXVI, and LXVII, the assignment of the absolute stereochemistry can be made to all of the intermediates, as indicated in Fig. 7.

The final steps in the pathway from LXXV or LXXVII to HTCA could involve the reverse elimination of water from these molecules to form *L-erythro*-(*E* or *Z*)-1,3,5,6-hex-1-enetetra-carboxylic acid and *L-threo*-(*E* or *Z*)-1,3,5,6-hex-1-enetetra-carboxylic acid, respectively. After equilibration of these two isomers by racemization at C-3, and specific reduction of the *L-erythro*-(*E* or *Z*)-1-HETCA, the final product HTCA would be produced. The occurrence of this racemization is confirmed by the rapid exchange of the C-3 hydrogens of *trans*-glutaconic acid and is a required step in the biosynthetic pathway because of the *meso* stereochemistry of the final HTCA (White *et al.*, 1996). This pathway is consistent with the experimental observation that incubation of cell extracts with *DL-arabino*-HHTCA leads to the production of *erythro*-(*E* or *Z*)-1-HETCA and HTCA.

Although this is a possible route for the final steps in the biosynthetic scheme, the experimental evidence would suggest that these last three reactions in fact occur with the coenzyme A derivative of LXXV and LXXVII. The experimental data show that the conversion of *DL-arabino*-HHTCA to *erythro*-(*E*)-1-HETCA and HTCA is stimulated by the addition of HSCoA and ATP to the incubation mixture. This reaction would be catalyzed by an enzyme(s) similar to the (*R*)-lactyl-CoA desaturase involved in the acrylic acid pathway of lactate metabolism (Kuchta and Abeles, 1985; Kuchta *et al.*, 1986; Hofmeister and Buckel, 1992) or to the (*R*)-2-hydroxyglutaryl-CoA dehydratase involved in the fermentation of glutamate via the hydroxyglutarate pathway (Klees *et al.*, 1992; Bendart *et al.*, 1993). All of these enzymes are oxygen sensitive, they all likely contain Fe/S centers, and they all catalyze the elimination of 2-hydroxyacids with the same (*R*) stereochemistry found in our intermediates. On the basis of the work in Buckel's laboratory (Buckel and Golding, 1999), reactions 17 and 18 appear to proceed by a radical mechanism with ketyl intermediates, as shown in Fig. 8. The *M. jannaschii* genome contains a gene for 2-hydroxyglutaryl-CoA dehydratase (MJ0007) and two genes for the activator protein of (*R*)-2-hydroxyglutaryl-CoA dehydratase (MJ0004 and MJ0800). An open reading frame predicted to code for the activator protein of (*R*)-2-hydroxyglutaryl-CoA dehydratase and the β subunit of (*R*)-2-hydroxyglutaryl-CoA dehydratase has also been identified in *Methanopyrus kandleri* (Vorholt *et al.*, 1997). We have now confirmed that the enzymes generated from the protein products of the *M. jannaschii* genes MJ0004 and MJ0007 function as a (*R*)-2-hydroxyglutaryl-CoA dehydratase and

are involved in the biosynthesis of β -glutamate. We thus propose that an enzyme composed of the gene products of both the MJ0007 and MJ0800 genes is involved in performing the same type of reaction on the intermediates LXXVIII and LXXIX.

Additional support for the involvement of the CoA ester in the final steps of HTCA biosynthesis comes from the fact that most of the known examples of the reduction of α,β -unsaturated acids occur as their CoA esters (Caldeira *et al.*, 1996; Sedlmeier and Simon, 1985; Weeks and Wakil, 1969). One advantage of using a CoA ester in these last steps of the reaction is that the carboxylate group of HTCA is activated at the correct position for its coupling to the α -amino group of glutamic acid in the biosynthesis of methanofuran.

In summary, we find that all of the reactions used in the construction of the HTCA molecule are analogous to reactions previously described in bacterial metabolism, and each reaction is likely catalyzed by paralogues of the enzymes catalyzing these reactions. This finding is analogous to that observed in the biosynthesis of coenzyme B.

V. BIOSYNTHESIS OF COENZYME F₄₂₀

Little work has been published on the biosynthesis of coenzyme F₄₂₀, a hydride transfer coenzyme that functions in a manner similar to NADH (Jacobson and Walsh, 1984). The most recent work published on F₄₂₀ biosynthesis is from Bacher's laboratory, where it has been demonstrated that growing cells of *M. thermoautotrophicum* readily incorporate 5-amino-6-ribitylamino-2,4-(1*H*,3*H*)-pyrimidinedione (XLIII) (Fig. 9) and 4-hydroxyphenylpyruvate (XLVI) into the 7,8-didemethyl-8-hydroxy-5-deazariboflavin moiety (LI) of coenzyme F₄₂₀ (CVIII) (Fo) (Reuke *et al.*, 1992). Compound XLIII, an established intermediate in riboflavin biosynthesis (Bacher *et al.*, 1996; Richter *et al.*, 1997), has been proposed to be formed by the same route as that found in bacterial riboflavin biosynthesis, and thus it represents the last common intermediate in the biosynthesis of riboflavin and Fo. The bacterial pathway, however, cannot be completely identical to the proposed riboflavin biosynthesis pathway in archaea since the methanoarchaea appear to lack GTP cyclohydrolase II, the first enzyme in riboflavin biosynthesis (Howell and White, 1997). Our current ideas for the pathway are outlined in Fig. 9, which shows a possible route for the formation of Fo (LI) from GTP, and in Fig. 10, which shows a possible route for the conversion of LI to coenzyme F₄₂₀. We have established, using incubations with cell extracts of *M. thermophila*, that XLIII and XLVI (Fig. 9) are the precursors for the formation of the Fo moiety of F₄₂₀ (LI), and that

325

3 COENZYME F₄₂₀

the biosynthesis of coenzyme F₄₂₀, functions in a manner similar to the most recent work published on laboratory, where it has been demonstrated that *M. thermoautotrophicum* readily incorporates 3H-pyrimidinone (XLIII) (Fig. 9) into the 7,8-didemethyl-8-hydroxy of coenzyme F₄₂₀ (CVIII) (Fo) II, an established intermediate in the pathway (Richter *et al.*, 1996; Richter *et al.*, 1997), has the same route as that found in bacteria. It represents the last common intermediate in the biosynthesis of riboflavin and Fo. The bacterial pathway is identical to the proposed riboflavin biosynthesis in the methanoarchaea appear to be the same enzyme in riboflavin biosynthesis. The ideas for the pathway are outlined in the text. The route for the formation of Fo (LI) is a possible route for the conversion of Fo (LI) to Fo (LI) as established, using incubations with Fo (LI) and XLIII (Fig. 9) are the Fo moiety of F₄₂₀ (LI), and that




FIG. 9. Proposed pathway for the biosynthesis of the Fo moiety of coenzyme F₄₂₀. **I**, Guanosine triphosphate; **II**, triaminopyrimidine triphosphate; **XL**, 2,5-diamino-6-ribosylamino-4(3*H*)-pyrimidinone 5'-phosphate; **XL1**, 2,5-diamino-6-ribosylamino-2,4(1*H*,3*H*)-pyrimidinedione 5'-phosphate; **XLII**, 5-amino-6-ribitylamino-2,4(1*H*,3*H*)-pyrimidinedione 5'-phosphate; **XLIII**, 5-amino-6-ribitylamino-2,4(1*H*,3*H*)-pyrimidinedione; **XLIV**, oxidized **XLIII**; **XLV**, 4-hydroxyphenylpyruvate; **XLVI**, enol form of 4-hydroxyphenylpyruvate; **XLVII**, condensation product between **XLIV** and **XLVI**; **XLVIII**, **XLIX**, and **L**, intermediates in the formation of Fo; **LI**, Fo, 7,8-dimethyl-8-hydroxy-5-deazariboflavin.

lactyl phosphate, formed by the GTP-dependent phosphorylation of lactate (CII), serves as an intermediate in the formation of the phosphodiester bond in F_{420} .

The first reaction in the methanogens is catalyzed by GTP cyclohydrolase III (Fig. 9, reaction 1). We propose that the triaminopyrimidine triphosphate (II) shown in Fig. 9 is the true intermediate in the archaeal pathway. This very reactive compound would therefore serve as the branch point for the biosynthesis of F_{420} , flavin adenine dinucleotide (FAD), methanopterin, and possibly also the 7-deazaguanosine-containing series of bases found in tRNA nucleosides (Watanabe *et al.*, 1997). After removal of pyrophosphate from II (Fig. 9, reaction 2), the resulting XL would be transformed to XLIII by the same series of reactions as occurs in riboflavin biosynthesis. Possible genes in the genome of *M. jannaschii* have been proposed to encode enzymes that catalyze the two reactions in the formation of XLIII (Koonin *et al.*, 1997). These are MJ0430 or MJ1102 for the *ribD* pyrimidine deaminase domain (Fig. 9, reaction 3) and MJ0671 for the *ribD* pyrimidine reductase domain (Fig. 9, reaction 4). We have now confirmed that MJ0671 is the gene encoding for the pyrimidine reductase and that the gene product of MJ0430 is a dCTP deaminase.

Compound XLIII, after a two-electron oxidation to compound XLIV, condenses with XLVI, likely as the enol XLV (Fig. 9, reaction 7), to form compound XLVII, which, after the series of indicated reactions, leads to the formation of reduced Fo. A proposed series of reactions is shown in Fig. 9. Reduced Fo would be readily oxidized to Fo or could proceed down the pathway as the reduced form of Fo (H_2Fo). The involvement of oxalate in the biosynthesis of Fo is supported by our demonstration of the presence of low concentrations of oxalate in cell extracts of *M. thermoautotrophicum* and our demonstration of oxalate production in a cell extract of *M. jannaschii* incubated with XLVI and XLIII. Predictions as to the identity of the enzyme(s) involved in the biosynthesis based on sequence comparisons are currently impossible since any analogous reactions for comparison are unknown. We considered it very likely that a single enzyme could catalyze reactions 7 to 11 in Fig. 9 based on similar complex transformations carried out by such enzymes as GTP cyclohydrolase I (Nar *et al.*, 1995) and 6,7-dimethyl-8-ribitylumazine synthase in riboflavin biosynthesis (Bacher *et al.*, 1996). This has now been confirmed in our laboratory, for an enzyme cloned from *M. jannaschii*.

We have also studied the formation of the phosphodiester bond in the coenzyme F_{420} structure and find that it is readily formed when cell extracts of *M. thermophila* or *M. jannaschii* are incubated with Fo (LI), GTP (I), and L-lactate (CII) (Fig. 10). The formation of this bond is great-

WHITE

-dependent phosphorylation of lactate in the formation of the phospho-

gens is catalyzed by GTP cyclohydrolase that the triaminopyrimidine is the true intermediate in the ar- compound would therefore serve as sis of F_{420} , flavin adenine dinu- possibly also the 7-deazaguanosine in tRNA nucleosides (Watanabe *et al.* phosphate from **II** (Fig. 9, reaction 2), led to **XLIII** by the same series of synthesis. Possible genes in the proposed to encode enzymes that formation of **XLIII** (Koonin *et al.*, for the *ribD* pyrimidine deaminase i71 for the *ribD* pyrimidine reduc- have now confirmed that MJ0671 line reductase and that the gene base.

ron oxidation to compound **XLIV**, ol **XLV** (Fig. 9, reaction 7), to form eries of indicated reactions, leads posed series of reactions is shown ly oxidized to F_0 or could proceed m of F_0 (H_2F_0). The involvement supported by our demonstration s of oxalate in cell extracts of *M.* stration of oxalate production in ted with **XLVI** and **XLIII**. Predict- e(s) involved in the biosynthesis currently impossible since any e unknown. We considered it very talyze reactions 7 to 11 in Fig. 9 ions carried out by such enzymes 1995) and 6,7-dimethyl-8-ribityl- nthesis (Bacher *et al.*, 1996). This atory, for an enzyme cloned from

of the phosphodiester bond in the it is readily formed when cell ex- *schii* are incubated with F_0 (**LI**), he formation of this bond is great-

METHANOGENIC COFACTORS

327




FIG. 10. Proposed pathway for the biosynthesis of coenzyme F_{420} from **LI**, **LI**, F_0 , 7,8-didemethyl-8-hydroxy-5-deazariboflavin; **CII**, L-lactate; **CIII**, L-lactyl phosphate; **CIV**, L-lactyl diphosphate; **CV**, L-lactyl triphosphate; **CVI**, F_{420} -0; **CVII**, F_{420} -1; **CVIII**, F_{420} -2, coenzyme F_{420} .

ly stimulated by including in the incubation mixture an unknown product obtained by treating L-lactic acid with $POCl_3$. At present, we propose the pathway shown in Fig. 10 as a possible pathway for the generation of F_{420} (F_{420} -2) and its γ -glutamyl derivatives (F_{420} -1, F_{420} -3, F_{420} -4,

etc). The first step in this pathway would begin with the GTP-dependent phosphorylation of L-lactate (CII) to L-lactyl-phosphate (CIII). The resulting CIII is then phosphorylated by GTP in two consecutive reactions, first to L-lactyldiphosphate (CIV) and then finally to L-lactyl triphosphate (CV). This series of reactions is the same that is used in nature in the biosynthesis of the phosphodiester in nucleic acids. The nature of the compound formed from treating L-lactic acid with POCl_3 , which stimulates CVI synthesis, has not as yet been identified, but it is known not to be CIV or CV. The current evidence indicates that it is a cyclic derivative of CIV that undergoes a ring opening to form CIV, which then serves as a precursor to CVI, as shown in Fig. 10.

We have confirmed the presence of lactoyl phosphate in *M. thermophila* and *M. thermoautotrophicum* ΔH by gas chromatography-mass spectrometry (GC-MS) analysis. We have also shown that cell extracts of *M. thermophila* incubated with [hydroxy- ^{18}O]lactate and GTP produce ^{18}O -labeled CIII.

Another important aspect of the coenzyme F_{420} structure is the presence of γ -linked glutamic acid residues. This structural element is also a common characteristic of some of the different methanofurans (Daniels, 1993) and folates, and it identifies another common structural element uniting these coenzymes, as does the pyrophosphate diester linkage present in NAD, FAD, coenzyme A, and molybdopterin. The mechanism and enzymology for the assembly of the γ -linked glutamic acid have been studied in bacteria and in eukaryotes, where the reaction is involved in the generation of the folyl- γ -polyglutamates (Banerjee *et al.*, 1988; Garrow *et al.*, 1992).

The enzymes responsible for these reactions are the folyl- γ -polyglutamate synthases (EC 6.3.2.17), and they have been found to be moderately conserved between bacteria and eukaryotes. A paper on the X-ray structure of the *Lactobacillus casei* enzyme reveals that the enzyme is a modular protein consisting of two domains, one with a typical mononucleotide-binding fold and the other strikingly like that in the folate-binding enzyme dihydrofolate reductase (Sun *et al.*, 1998). If the archaea contained an enzyme(s) related to this enzyme, we would expect at least a portion of the enzyme sequence to have some homology to the folylpolyglutamate synthases. However, no open reading frames in the genomes of *M. jannaschii* or *M. thermoautotrophicum* ΔH can be found that have sequences homologous to any known folylpolyglutamate synthase. We have three possible explanations for this finding: (1) the sequences of the archaeal enzymes have diverged to such an extent from the known enzymes that it is impossible to detect them by sequence homologies, (2) a nonortholog replacement for these enzymes

I. WHITE

ould begin with the GTP-dependent
L-lactyl phosphate (CIII). The re-
l by GTP in two consecutive reac-
CIV) and then finally to L-lactyl
actions is the same that is used in
osphodiester in nucleic acids. The
treating L-lactic acid with POCl_3 ,
not as yet been identified, but it is
rent evidence indicates that it is a
goes a ring opening to form CIV,
CVI, as shown in Fig. 10.

of lactoyl phosphate in *M. ther-*
um ΔH by gas chromatography-
s. We have also shown that cell ex-
with [^3H]lactate and GTP

coenzyme F_{420} structure is the
residues. This structural element
me of the different methanofurans
antifies another common structur-
as does the pyrophosphate diester
zyme A, and molybdopterin. The
assembly of the γ -linked glutamic
nd in eukaryotes, where the reac-
he folyl- γ -polyglutamates (Baner-

reactions are the folyl- γ -polyglu-
they have been found to be mod-
nd eukaryotes. A paper on the X-
zi enzyme reveals that the enzyme
two domains, one with a typical
other strikingly like that in the fo-
ductase (Sun *et al.*, 1998). If the
ted to this enzyme, we would ex-
sequence to have some homology
however, no open reading frames in
thermoautotrophicum ΔH can be
s to any known folypolyglutamate
anations for this finding: (1) the
have diverged to such an extent
impossible to detect them by se-
; replacement for these enzymes

METHANOGENIC COFACTORS

329




FIG. 11. Proposed pathway for the biosynthesis of coenzyme M. CX, Phosphoenol pyruvate; CXI, bisulfite; CXII, L-sulfolactyl phosphate; CXIII, L-sulfolactate; CXIV, sulfolpyruvate; CXV, sulfoacetaldehyde; CXVI, 2-(sulfomethyl)thiazolidine-4-carboxylic acid; CXVII, sulfoethylcysteine; CXVIII, coenzyme M, 2-mercaptoethanesulfonic acid.

exists in the archaea, or (3) the peptide bond-forming reaction proceeds by another mechanism and would thus be expected to use a different enzyme. We have also identified an α -glutamyl-linked F_{420} in *M. jannaschii* that further establishes biochemical similarities between bacterial metabolism, which also produces α -glutamyl-linked folates (Ferone *et al.*, 1986), and the metabolism found in the Archaea.

VI. BIOSYNTHESIS OF COENZYME M

The currently established pathway for the biosynthesis of coenzyme M is outlined in Fig. 11. We have established that reaction 3 is carried out by the gene product of the *M. jannaschii* gene MJ1425 (Graupner and White, 2000). The TPP-dependent decarboxylation of reaction 4 is carried out by the enzyme composed from the two proteins produced by the MJ0256 gene, which are designated as ComD and ComE. These correspond to the MTH1206 and MTH1207 genes in *M. thermoautotrophicum* Δ H. The ComD and ComE proteins make an $\alpha_6\beta_6$ dodecamer enzyme that is a member of the acetolactate and pyruvate oxidase family of enzymes (Chang, 1992). Both subunits align with different portions of the phosphopyruvate decarboxylase (Schwartz *et al.*, 1998), an enzyme involved in the biosynthesis of compounds containing a P—C bond (Hidaka *et al.*, 1995).

We are currently trying to establish the identity of the other genes in the pathway, each of which has an analogous reaction occurring in other known enzymatic reactions. These include formation of the sulfonic acid derivative by a route analogous to that involved in the biosynthesis of UDP-sulfoquinovose, an intermediate in the biosynthesis of sulfolipids (Fig. 11, reaction 1) (Benning, 1998); the Stickland reaction used for reduction of the C—N bond in the thiazolidine intermediate (reaction 5) that occurs in D-proline reductase (Kabisch *et al.*, 1999); and a β -elimination of pyruvate and ammonia from sulfoethylcysteine (reaction 7) by a reaction like that catalyzed by β -cystathionase (Flavin and Slaughter, 1964).

REFERENCES

- Aurora, R., and Rose, G. D. (1998). Seeking an ancient enzyme in *Methanococcus jannaschii* using ORF, a program based on predicted secondary structure comparisons. *Proc. Natl. Acad. Sci. U.S.A.* **95**, 2818–2823.
- Bacher, A., Eberhart, S., and Richter, G. (1996). Biosynthesis of riboflavin. In "Escherichia coli and Salmonella typhimurium: Cellular and Molecular Biology" (F. C. Neidhardt *et al.*, eds.), 2nd ed., Vol. 1, pp. 657–664. American Society for Microbiology, Washington, DC.
- Banerjee, R. V., Shane, B., McGuire, J. J., and Coward, J. K. (1988). Dihydrofolate syn-

WHITE

METHANOGENIC COFACTORS

331

the bond-forming reaction proceeds as is expected to use a different α -glutamyl-linked F_{420} in *M. jannaschii* similarities between bacterial α -glutamyl-linked folates (Ferredoxin) found in the Archaea.

F COENZYME M

for the biosynthesis of coenzyme M. It has been established that reaction 3 is carried out by the *naschii* gene MJ1425 (Graupner et al., 1998). The decarboxylation of reaction 4 is carried out by the two proteins produced by the *comD* and *comE* genes. These correspond to the *comD* and *comE* genes in *M. thermoautotrophicus*. These two proteins make an $\alpha_6\beta_6$ dodecamer enzyme and pyruvate oxidase family. The amino acid sequences align with different portions of the *comD* and *comE* genes (Schwartz et al., 1998), an enzyme containing a P—C bond.

the identity of the other genes in the biosynthetic pathway. The reaction occurring in the biosynthesis of the sulfonic acid that involved in the biosynthesis of the sulfonic acid in the biosynthesis of sulfolipids (1998); the Stickland reaction in the thiazolidine intermediate (Kabisch et al., 1999); and the reaction from sulfoethylcysteine (reduced by β -cystathionase (Flavin

CES

the ancient enzyme in *Methanococcus jannaschii* predicted secondary structure comparisons.

36). Biosynthesis of riboflavin. In "Essentials of Cellular and Molecular Biology" (F. C. Campbell, ed.), pp. 57–664. American Society for Microbiology, Washington, D.C.

Coward, J. K. (1988). Dihydrofolate syn-

thase and folylpolyglutamate synthetase: Direct evidence for intervention of acyl phosphate intermediates. *Biochemistry* 27, 9062–9070.

Beinert, H., Kennedy, M. C., and Stout, C. D. (1996). Aconitase as iron-sulfur protein, enzyme, and iron-regulatory protein. *Chem. Rev.* 96, 2335–2373.

Bendart, K., Müller, U., Klees, A.-G., and Buckel, W. (1993). Identification of the gene encoding the activator of (R)-2-hydroxyglutaryl-CoA dehydratase from *Acidaminococcus fermentans* by gene expression in *Escherichia coli*. *FEBS Lett.* 329, 329–331.

Benning, C. (1998). Biosynthesis and function of the sulfolipids sulfoquinovosyl diacylglycerol. *Annu. Rev. Physiol. Mol. Biol.* 49, 53–76.

Bock, A.-K., Kunow, J., Glasemacher, J., and Schönheit, P. (1996). Catalytic properties, molecular composition and sequence alignments of pyruvate: Ferredoxin oxidoreductase from the methanogenic archaeon *Methanosarcina barkeri* (strain Fusaro). *Eur. J. Biochem.* 237, 35–44.

Buckel, W., and Golding, T. B. (1999). Radical species in the catalytic pathways of enzymes from anaerobes. *FEMS Microbiol. Rev.* 22, 523–541.

Bult, C. J., White, O., Olsen, G. J., Zhou, L., Fleischmann, R. D., Sutton, G. G., Blake, J. A., FitzGerald, L. M., Clayton, R. A., Gocayne, J. D., Kerlavage, A. R., Dougherty, B. A., Tomb, J.-F., Adams, M. D., Reich, C. I., Overbeek, R., Kirkness, E. F., Weinstock, K. G., Merrick, J. M., Glodek, A., Scott, J. L., Geoghegan, N. S. M., Weidman, J. F., Fuhrmann, J. L., Nguyen, D., Utterback, T. R., Kelley, J. M., Peterson, J. D., Sadow, P. W., Hanna, M. C., Cotton, M. D., Roberts, K. M., Hurst, M. A., Kaine, B. P., Borodovsky, M., Klenk, H.-P., Fraser, C. M., Smith, H. O., Woese, C. R., and Venter, J. C. (1997). Complete genome sequence of the methanogenic archaeon, *Methanococcus jannaschii*. *Science* 273, 1058–1073.

Caldeira, J., Feicht, R., White, H., Teixeira, M., Moura, J. J. G., Simon, H., and Moura, I. (1996). EPR and Mössbauer spectroscopic studies on enolase reductase. *J. Biol. Chem.* 271, 18743–18748.

Carreras, C. W., and Santi, D. V. (1995). The catalytic mechanism and structure of thymidylate synthase. *Annu. Rev. Biochem.* 64, 721–762.

Chang, Y. Y. (1992). Common ancestry of *Escherichia coli* pyruvate oxidase and the acetylhydroxy acid synthase of the branched-chain amino acid biosynthetic pathway. In "The Evolution of Metabolic Function" (R. P. Mortlock, ed.), pp. 81–104. CRC Press, Boca Raton, FL.

Christoserdova, L., Vorholt, J. A., Thauer, R. K., and Lidstrom, M. E. (1998). C-1 transfer enzymes and coenzymes linking methylotrophic bacteria and methanogenic archaea. *Science* 281, 99–102.

Daniels, L. (1993). Biochemistry of methanogens. In "The Biochemistry of the Archaea (Archaeobacteria)" (M. Kates, D. J. Kushner, and A. T. Matheson, eds.), pp. 41–112. Elsevier, Amsterdam.

DiMarco, A. A., Bobik, T. A., and Wolfe, R. S. (1990). Unusual coenzymes of methanogenesis. *Annu. Rev. Biochem.* 59, 355–394.

Eck, R., and Simon, H. (1994a). Preparation of (S)-2-substituted succinates by stereospecific reductions of fumarate and derivatives with resting cells of *Clostridium formicoaceticum*. *Tetrahedron* 50, 13631–13640.

Eck, R., and Simon, H. (1994b). Preparation of both enantiomers of malic and citramalic acid and other hydroxysuccinic acid derivatives by stereospecific hydrations of cis- or trans-2-butene-1,4-dioic acids with resting cells of *Clostridium formicoaceticum*. *Tetrahedron* 50, 13641–13654.

Eggen, R. I. L., Geerling, A. C. M., Roshoven, A. B. P., and de Vos, W. M. (1991). Cloning, sequence analysis, and functional expression of the acetyl coenzyme A synthase gene from *Methanoxanthus soehngenii*. *J. Bacteriol.* 173, 6383–6389.

- Eisenreich, W., and Bachor, A. (1992). Biosynthesis of methanofuran in *Methanobacterium thermoautotrophicum*. *J. Biol. Chem.* 267, 17574-17580.
- Ferone, R., Hanlon, M., Hunt, D., and Singer, S. (1986). Carboxyl-linked glutamates in the folylpolyglutamates of *Escherichia coli*. *J. Biol. Chem.* 261, 16356-16362.
- Flavin, M., and Slaughter, C. (1964). Cystathionine cleavage enzymes of *Neurospora*. *J. Biol. Chem.* 239, 2212-2219.
- Flint, D. H., and Allen, R. M. (1996). Iron-sulfur proteins with nonredox functions. *Chem. Rev.* 96, 2315-2334.
- Foor, F., and Brown, G. M. (1975). Purification and properties of guanosine triphosphate cyclohydrolase from *Escherichia coli*. *J. Biol. Chem.* 250, 3545-3551.
- Garrow, T. A., Admon, A., and Shane, B. (1992). Expression cloning of a human cDNA encoding folylpoly(γ -glutamate) synthetase and determination of its primary structure. *Proc. Natl. Acad. Sci. U.S.A.* 89, 9151-9155.
- Gawron, O., and Mahajan, K. P. (1966). α -Methyl-cis-aconitic acid. Aconitase substrate. II. Substrate properties and aconitase mechanism. *Biochemistry* 5, 2343-2350.
- Gehring, U., and Arnon, D. I. (1972). Purification and properties of α -ketoglutarate synthase from a photosynthetic bacterium. *J. Biol. Chem.* 247, 6963-6969.
- Glasemacher, J., Bock, A.-K., Schmid, R., and Schönheit, P. (1997). Purification and properties of acetyl-CoA synthetase (ADP-forming), an archael enzyme of acetate formation and ATP synthesis, from the hyperthermophilic *Pyrococcus furiosus*. *Eur. J. Biochem.* 244, 561-567.
- Gorkovenko, A., Roberts, M. F., and White, R. H. (1994). Identification, biosynthesis and function of 1,3,4,6-hexanetetra-carboxylic acid in *Methanobacterium thermoautotrophicum* strain Δ H. *Appl. Environ. Microbiol.* 60, 1249-1253.
- Graupner, M., and White, R. H. (2000). Analysis of the coenzymes present in *Methanococcus jannaschii*. *J. Bacteriol.* (in preparation).
- Graupner, M., Xu, H., and White, R. H. (2000). Identification of an Archaeal 2-hydroxy acid dehydrogenase involved in coenzyme biosynthesis in Methanarchaea. *J. Bacteriol.* 182, 3688-3692.
- Green, J. M., Nichol, B. P., and Matthews, R. G. (1996). Folate biosynthesis, reduction, and polyglutamylation. In *"Escherichia coli and Salmonella typhimurium: Cellular and Molecular Biology"* (F. C. Neidhardt et al., eds.), 2nd ed., Vol. 1, pp. 665-673. American Society for Microbiology, Washington, DC.
- Heider, J., Mai, X., and Adams, M. W. W. (1996). Characterization of 2-ketoisovalerate ferredoxin oxidoreductase, a new reversible coenzyme A-dependent enzyme involved in peptide fermentation by hyperthermophilic archaea. *J. Bacteriol.* 178, 780-787.
- Heim, S., Kunkel, A., Thauer, R. K., and Hedderich, R. (1998). Thiol:fumarate reductase (Tfr) from *Methanobacterium thermoautotrophicum*. Identification of the catalytic sites for fumarate reduction and thiol oxidation. *Eur. J. Biochem.* 253, 292-299.
- Hennig, M., D'Arcy, A., Hampele, I., Page, M. G. P., Oefner, C., and Dale, G. E. (1998). Crystal structure and reaction mechanism of 7,8-dihydropterin aldolase from *Staphylococcus aureus*. *Nat. Struct. Biol.* 5, 357-362.
- Hennig, M., Dale, G. E., D'Arcy, A., Danel, F., Fischer, S., Gray, C. P., Jolidon, S., Muller, F., Page, M. G. P., Pattison, P., and Oefner, C. (1999). The structure and function of the 6-hydroxymethyl-7,8-dihydropterin pyrophosphokinase from *Haemophilus influenzae*. *J. Mol. Biol.* 287, 211-219.
- Hidaka, T., Goda, M., Kuzuyama, T., Takei, N., Hidaka, M., and Seto, H. (1995). Cloning and nucleotide sequence of fosfomycin biosynthetic genes of *Streptomyces wedmorensis*. *Mol. Gen. Genet.* 249, 274-280.
- Hofmeister, A. E. M., and Buckel, W. (1992). (R)-Lactyl-CoA dehydratase from *Clostridi-*

WHITE

METHANOGENIC COFACTORS

333

- thesis of methanofuran in *Methanobacterium*. *J. Biol. Chem.* **267**, 17574-17580.
- , S. (1986). Carboxyl-linked glutamates in *li*. *J. Biol. Chem.* **261**, 16356-16362.
- imine cleavage enzymes of *Neurospora*. *J. Biol. Chem.* **267**, 17574-17580.
- ir proteins with nonredox functions. *Chem. Rev.* **75**, 1-10.
- and properties of guanosine triphosphate. *J. Biol. Chem.* **250**, 3545-3551.
- l. Expression cloning of a human cDNA encoding a protein with a structure similar to that of the primary structure of the *li* protein. *J. Biol. Chem.* **267**, 17574-17580.
- thyl-cis-aconitic acid. Aconitase substrate. *J. Biol. Chem.* **267**, 17574-17580.
- chanism. *Biochemistry* **5**, 2343-2350.
- ion and properties of α -ketoglutarate synthase. *J. Biol. Chem.* **267**, 17574-17580.
- Schönheit, P. (1997). Purification and properties of an archaeal enzyme of acetate-forming activity from the hyperthermophile *Pyrococcus furiosus*. *Eur. J. Biochem.* **247**, 6963-6969.
- H. (1994). Identification, biosynthesis and function of the coenzyme B in *Methanobacterium thermoautotrophicum*. *J. Biol. Chem.* **269**, 1249-1253.
- is of the coenzymes present in *Methanobacterium thermoautotrophicum*. *J. Biol. Chem.* **269**, 1249-1253.
- l). Identification of an Archaeal 2-hydroxy-3-oxo-4-oxopentanoate synthase in *Methanobacterium thermoautotrophicum*. *J. Biol. Chem.* **269**, 1249-1253.
- G. (1996). Polate biosynthesis, reduction, and function in *Salmonella typhimurium*: Cellular localization and function. *J. Biol. Chem.* **271**, 665-673.
- ngton, DC.
- 96). Characterization of 2-ketoisovalerate decarboxylase, a coenzyme A-dependent enzyme involved in the biosynthesis of coenzyme B in the hyperthermophilic archaea. *J. Bacteriol.* **178**, 780-787.
- erich, R. (1998). Thiol:fumarate reductase from *Methanobacterium thermoautotrophicum*. Identification of the catalytic site. *Eur. J. Biochem.* **253**, 292-299.
- . G. P., Oefner, C., and Dale, G. E. (1998). Identification of the catalytic site of 7,8-dihydropyrimidin aldolase from *Methanobacterium thermoautotrophicum*. *J. Biol. Chem.* **273**, 357-362.
- Fischer, S., Gray, C. P., Jolidon, S., Muller, R., and C. (1999). The structure and function of pyrophosphokinase from *Haemophilus influenzae*. *J. Biol. Chem.* **274**, 1249-1253.
- ., Hidaka, M., and Seto, H. (1995). Cloning and characterization of the gene for the biosynthetic enzyme of coenzyme B in *Methanobacterium thermoautotrophicum*. *J. Biol. Chem.* **270**, 1249-1253.
- R)-Lactyl-CoA dehydratase from *Clostridium propionicum*. Stereochemistry of the dehydration of (R)-2-hydroxybutyryl-CoA to crotonyl-CoA. *Eur. J. Biochem.* **206**, 547-552.
- Howell, D. M., and White, R. H. (1997). D-erythro-Neopterin biosynthesis in the methanogenic archaea *Methanococcus thermophilus* and *Methanobacterium thermoautotrophicum* Δ II. *J. Bacteriol.* **179**, 5165-5170.
- Howell, D. M., Harich, K., Xu, H., and White, R. H. (1998). The α -keto acid chain elongation reactions involved in the biosynthesis of coenzyme B (7-mercaptoheptanoylthreonine phosphate) in methanogenic archaea. *Biochemistry* **37**, 10108-10117.
- Howell, D. M., Graupner, M., and White, R. H. (2000a). Biosynthesis of the 1,3,4,6-hexanetetra-carboxylic acid moiety of methanofuran in the methanobacteria. *J. Am. Chem. Soc.* (submitted for publication).
- Howell, D. M., Graupner, M., Xu, H., and White, R. H. (2000b). Identification of enzymes homologous to isocitrate dehydrogenase that are involved in coenzyme B and leucine biosynthesis in the methanobacteria. *J. Bacteriol.* (in press).
- Jacob, U., Mack, M., Clausen, T., Huber, R., Buckel, W., and Messerschmidt, A. (1997). Glutamate CoA-transferase from *Acidaminococcus fermentans*: The crystal structure reveals homology with other CoA-transferases. *Structure* **5**, 415-426.
- Jacobson, F., and Walsh, C. (1984). Properties of 7,8-didemethyl-8-hydroxy-5-deazaflavins relevant to redox coenzyme function in methanogen metabolism. *Biochemistry* **23**, 979-988.
- Jetten, M. S. M., Stams, A. J. M., and Zehnder, A. J. B. (1989). Isolation and characterization of acetyl-coenzyme A synthetase from *Methanothermobacter thermoautotrophicus*. *J. Bacteriol.* **171**, 5430-5435.
- Kabisch, U. C., Grätzdörffler, A., Schierhorn, A., Rücknagel, K. P., Andreesen, J. R., and Pich, A. (1999). Identification of D-proline reductase from *Clostridium stricklandii* as a selenoenzyme and indications for a catalytically active pyruvoyl group derived from a cysteine residue by cleavage of a propeptide. *J. Biol. Chem.* **274**, 8445-8454.
- Kates, M. (1993). Membrane lipids of archaea. In "The Biochemistry of Archaea (Archaeobacteria)" (M. Kates, D. J. Kushner, and A. T. Matheson, eds.), pp. 261-295. Elsevier, Amsterdam.
- Kim, M.-J., and Whitesides, G. M. (1988). L-Lactate dehydrogenase: Substrate specificity and use as a catalyst in synthesis of homochiral 2-hydroxy acids. *J. Am. Chem. Soc.* **110**, 2959-2964.
- Kloos, A.-G., Linder, D., and Buckel, W. (1992). 2-Hydroxyglutaryl-CoA dehydratase in *Fusobacterium nucleatum* (subsp. *nucleatum*): An iron-sulfur flavoprotein. *Arch. Microbiol.* **158**, 294-301.
- Klenk, H. P., Clayton, R. A., Tomb, J. F., White, O., Nelson, K. E., Ketchum, K. A., Dodson, R. J., Gwinn, M., Hickey, E. K., Peterson, J. D., Richardson, D. L., Kerlavage, A. R., Graham, D. E., Kyrpides, N. C., Fleischmann, R. D., Quackenbush, J., Lee, N. H., Sutton, G. G., Gill, S., Kirkness, E. F., Dougherty, B. A., McKenney, K., Adams, M. D., Loftus, B., Peterson, S., Reich, C. I., McNeil, L. K., Badger, J. H., Glodek, A., Zhou, L. X., Overbeck, R., Gocayne, J. D., Weidman, J. F., McDonald, L., Utterback, T., Cotton, M. D., Spriggs, T., Artiach, P., Kaine, B. P., Sykes, S. M., Sadow, P. W., D'Andrea, K. P., Bowman, C., Fujii, C., Garland, S. A., Mason, T. M., Olsen, G. J., Fraser, C. M., Smith, H. O., Woese, C. R., and Venter, J. C. (1998). The complete genome sequence of the hyperthermophilic, sulfate-reducing archaeon *Archaeoglobus fulgidus*. *Nature (London)* **390**, 364-370.
- Kletzin, A., and Adams, M. W. W. (1996). Molecular and phylogenetic characterization of pyruvate and 2-ketoisovalerate ferredoxin oxidoreductases from *Pyrococcus furiosus*.

- and pyruvate ferredoxin oxidoreductase from *Thermotoga maritima*. *J. Bacteriol.* 178, 248-257.
- Klinman, J. P., and Rose, I. A. (1971). Purification and kinetic properties of aconitase isomerase from *Pseudomonas putida*. *Biochemistry* 10, 2253-2266.
- Koonin, E. V., Mushengian, A. R., Galperin, M. Y., and Walker, D. R. (1997). Comparison of archaeal and bacterial genomes: Computer analysis of protein sequences predicts novel functions and suggests a chimeric origin for the archaea. *Mol. Microbiol.* 25, 619-637.
- Kuchta, R. D., and Abeles, R. H. (1985). Lactate reduction in *Clostridium propionicum*, purification and properties of lactyl-CoA dehydratase. *J. Biol. Chem.* 260, 13181-13189.
- Kuchta, R. D., Hanson, G. R., Holmquist, B., and Abeles, R. H. (1986). Fe-S centers in lactyl-CoA dehydratase. *Biochemistry* 25, 7301-7307.
- Kunitani, M. G., and Santi, D. V. (1980). On the mechanism of the 2'-deoxyuridylate hydroxymethylase. *Biochemistry* 19, 1271-1275.
- Latimer, M. T., and Ferry, J. G. (1993). Cloning, sequence analysis, and hyperexpression of the genes encoding phosphotransacetylase and acetate kinase from *Methanosarcina thermophila*. *J. Bacteriol.* 175, 6822-6829.
- Leigh, J. A. (1983). Levels of water-soluble vitamins in methanogenic and non-methanogenic bacteria. *Appl. Environ. Microbiol.* 45, 800-803.
- Lopez, P., and Lacks, S. A. (1993). A bifunctional protein in the folate biosynthetic pathway of *Streptococcus pneumoniae* with dihydroneopterin aldolase and hydroxymethyl-dihydropterin pyrophosphokinase activities. *J. Bacteriol.* 175, 2214-2220.
- Matthews, R. G. (1996). One-carbon metabolism. In *Escherichia coli and Salmonella typhimurium: Cellular and Molecular Biology* (F. C. Neidhardt et al., eds.), 2nd ed., Vol. 1, pp. 600-611. American Society for Microbiology, Washington, DC.
- Menon, S., and Ragsdale, S. W. (1997). Mechanism of the *Clostridium thermoaceticum* pyruvate:ferredoxin oxidoreductase: Evidence for the common catalytic intermediacy of the hydroxyethylthiamine pyrophosphate radical. *Biochemistry* 36, 8484-8494.
- Murry, B. W., Wittmann, V., Burkart, M. D., Hung, S.-C., and Wong, C.-H. (1997). Mechanism of human α -1,3-fucosyltransferase V: Glycosidic cleavage occurs prior to nucleophilic attack. *Biochemistry* 36, 823-831.
- Musick, W. D. L. (1981). Structural features of the phosphoribosyltransferases and their relationships to the human deficiency disorder of purine and pyrimidine metabolism. *Crit. Rev. Biochem.* 11, 1-34.
- Nar, H., Huber, R., Auerbach, G., Fischer, M., Hosl, C., Ritz, H., Bacher, A., Meining, W., Eberhart, S., and Bacher, A. (1995). Active site topology and reaction mechanism of GTP cyclohydrolase I. *Proc. Natl. Acad. Sci. U.S.A.* 92, 12120-12125.
- Neilson, N. E. (1956a). The presence of aconitase and "aconitic hydratase" in *Aspergillus niger*. *J. Bacteriol.* 71, 356-361.
- Neilson, N. E. (1956b). Aconitic hydratase in *Aspergillus niger*. In *Methods in Enzymology* (S. P. Colowick and N. O. Kaplan, eds.), Vol. 5, pp. 614-616. Academic Press, New York.
- Neumann, S., and Simon, H. (1984). On a non-pyridine nucleotide-dependent 2-oxo-acid reductase of broad substrate specificity from two *Proteus* species. *FEBS Lett.* 167, 29-32.
- Pomper, B. K., Vorholt, J. A., Chistoserdova, L., Lidstrom, M. E., and Thauer, R. K. (1999). A methenyl tetrahydromethanopterin cyclohydrolase and a methenyl tetrahydrofolate cyclohydrolase in *Methylobacterium extorquens* AM1. *Eur. J. Biochem.* 261, 475-480.
- Rasche, M. E., and White, R. H. (1998). Mechanism of the enzymatic formation of 4-(β -D-

WHITE

METHANOGENIC COFACTORS

335

from *Thermotoga maritima*. *J. Bacteriol.*

on and kinetic properties of aconitate isomerase. *Biochemistry* 10, 2253-2266.

Y., and Walker, D. R. (1997). Comparison of protein sequences predicts a common origin for the archaea. *Mol. Microbiol.* 25,

reduction in *Clostridium propionicum*, aconitase. *J. Biol. Chem.* 260, 13181-

and Abeles, R. H. (1986). Fe-S centers in aconitase. *J. Biol. Chem.* 261, 7301-7307.

the mechanism of the 2'-deoxyuridylate hydroxymethyltransferase.

sequence analysis, and hyperexpression of aconitase and acetate kinase from *Methanosarcina*.

vitamins in methanogenic and non-methanogenic bacteria. *Microbiol. Rev.* 45, 800-803.

protein in the folate biosynthetic pathway. *J. Bacteriol.* 175, 2214-2220.

in "Escherichia coli and Salmonella typhimurium" (F. C. Neidhardt et al., eds.), 2nd ed., ASM Press, Washington, DC.

ism of the *Clostridium thermoaceticum* aconitase for the common catalytic intermediate radical. *Biochemistry* 36, 8484-8494.

ng, S.-C., and Wong, C.-H. (1997). Mechanism of aconitase: Glycosidic cleavage occurs prior to nucleophilic attack of the phosphoribosyltransferase and their role in purine and pyrimidine metabolism.

sl, C., Ritz, H., Bacher, A., Meining, W., site topology and reaction mechanism of aconitase. *J. Biol. Chem.* 267, 12120-12125.

se and "aconitic hydratase" in *Aspergillus*

niger. In "Methods in Enzymology", Vol. 5, pp. 614-616. Academic Press, New York.

uridine nucleotide-dependent 2-oxo-acid aconitase in two *Proteus* species. *FEBS Lett.* 167,

idstrom, M. E., and Thauer, R. K. (1999). Aconitase and a methenyl tetrahydrofolate synthetase. *Eur. J. Biochem.* 261, 475-480.

ism of the enzymatic formation of 4-β-D-

ribofuranosyl)aminobenzene 5'-phosphate during the biosynthesis of methanopterin. *Biochemistry* 37, 11343-11351.

Reuke, B., Korn, S., Eisenreich, W., and Bacher, A. (1992). Biosynthetic precursors of deazaflavins. *J. Bacteriol.* 174, 4042-4049.

Richter, G., Fischer, M., Krieger, C., Eberhart, S., Lütgen, H., Gerstenschlager, I., and Bacher, A. (1997). Biosynthesis of riboflavin: Characterization of the bifunctional deaminase-reductase in *Escherichia coli* and *Bacillus subtilis*. *J. Bacteriol.* 179, 2022-2028.

Rodwell, V. W. (1969). In "Metabolic Pathways" (D. M. Greenberg, ed.), 3rd ed., Vol. 3, pp. 317-373. Academic Press, New York.

Schwartz, D., Recktenwald, J., Pelzer, S., and Wohlleben, W. (1998). Isolation and characterization of the PEP-phosphomutase and the phosphonopyruvate decarboxylase genes from the phosphinothricin tripeptide producer *Streptomyces viridochromogenes* Tu494. *FEMS Microbiol. Lett.* 163, 149-157.

Sedlmayr, H., and Simon, H. (1985). Purification and some properties of an acryloyl-CoA reductase of *Clostridium kluyveri*. *Biol. Chem. Hoppe-Seyler* 366, 953-961.

Siebert, G., Carsotis, M., and Plaut, G. W. E. (1957). The enzymatic properties of isocitric dehydrogenase. *J. Biol. Chem.* 226, 977-991.

Simon, H., Hader, J., Günther, H., Neumann, S., and Thanos, J. (1985). Chiral compounds synthesized by biocatalytic reductions. *Angew. Chem., Int. Ed. Engl.* 24, 539-553.

Smith, D. R., Doucette-Stamm, L. A., Deloughery, C., Lee, H., Dubois, J., Aldredge, T., Bashirzadeh, R., Blakely, D., Cook, R., Gilbert, K., Harrison, D., Hoang, L., Keagle, P., Lamm, W., Pothier, B., Qiu, D., Spadafora, R., Vicaire, R., Wang, Y., Wierzbowski, J., Gibson, R., Jiwani, N., Caruso, A., Bush, D., Safer, H., Patwell, D., Prabhakar, S., McDougall, S., Shimer, G., Goyal, A., Pietrovski, S., Church, G. M., Daniels, C. J., Mao, J., Rice, P., Nölling, J., and Reeve, J. N. (1997). Complete genome sequence of *Methanobacterium thermoautotrophicum* ΔH: Functional analysis and comparative genomics. *J. Bacteriol.* 179, 7135-7156.

Solow, B. T., and White, R. H. (1997a). Absolute stereochemistry of 2-hydroxyglutaric acid present in methanopterin. *Chirality* 9, 678-680.

Solow, B. T., and White, R. H. (1997b). Biosynthesis of the peptide bond in the coenzyme N-(7-mercaptoheptanoyl)-L-threonine phosphate. *Arch. Biochem. Biophys.* 345, 299-304.

Stammers, D. K., Achari, A., Somers, D. O., Bryant, P. K., Rosemond, J., Scott, D. L., and Champness, J. N. (1999). 2.0 Å X-ray structure of the ternary complex of 7,8-dihydro-6-hydroxymethylpterinpyrophosphokinase from *Escherichia coli* with ATP and a substrate analogue. *FEBS Lett.* 456, 49-53.

Sun, X. L., Bogner, A. L., Baker, E. N., and Smith, C. A. (1998). Structural homologies with ATP- and folate-binding enzymes in the crystal structure of fulylpolylglutamate synthetase. *Proc. Natl. Acad. Sci. U.S.A.* 95, 6647-6652.

Talarico, T. L., Ray, P. H., Dev, I. K., Merrill, B. M., and Dallas, W. S. (1992). Cloning, sequence analysis, and overexpression of *Escherichia coli* folK, the gene coding for 7,8-dihydro-6-hydroxymethylpterin-pyrophosphate. *J. Bacteriol.* 174, 5971-5977.

Telen, T. T., and van Milligan, H. (1960). The stereochemistry of the succinic dehydrogenase reaction. *J. Am. Chem. Soc.* 82, 4115-4116.

Tersteegen, A., Linder, D., Thauer, R. K., and Hedderich, R. (1997). Structures and functions of four anabolic 2-oxoacid oxidoreductases in *Methanobacterium thermoautotrophicum*. *Eur. J. Biochem.* 244, 862-868.

Thauer, R. K. (1998). Biochemistry of methanogenesis: A tribute to Marjory Stephenson. *Microbiology (Amsterdam)* 144, 2377-2406.

- Thompson, H., Tersteegen, A., Thauer, R. K., and Hedderich, R. (1998). Two malate dehydrogenase in *Methanobacterium thermoautotrophicum*. *Arch. Microbiol.* 170, 38-42.
- van Hellemond, J. J., and Tielens, A. G. M. (1994). Expression and functional properties of fumarate reductase. *Biochem. J.* 304, 321-331.
- Vorholt, J. A., Vaupel, M., and Thauer, R. T. (1997). A selenium-dependent and a selenium-independent formylmethanofuran dehydrogenase and their transcriptional regulation in the hyperthermophilic *Methanopyrus kandleri*. *Mol. Microbiol.* 23, 1033-1042.
- Wang, C. C., and Barker, H. A. (1969). Purification and properties of L-citramalate hydrolyase. *J. Biol. Chem.* 244, 2516-2526.
- Watanabe, M., Matsuo, M., Tanaka, S., Akimoto, H., Asahi, S., Nishimura, S., Katze, J. R., Hashizume, T., Crain, P. F., McCloskey, J. A., and Okada, N. (1997). Biosynthesis of archaeosine, a novel derivative of 7-deazaguanosine specific to archaeal tRNA, proceeds via a pathway involving base replacement on the tRNA polynucleotide chain. *J. Biol. Chem.* 272, 20146-20151.
- Weeks, G., and Wakil, S. J. (1969). Enoyl acyl carrier protein reductases from *Escherichia coli*. In "Methods in Enzymology" (J. M. Lowenstein, ed.), Vol. 14, pp. 66-73. Academic Press, New York.
- Weiss, D. S., and Thauer, R. K. (1993). Methanogenesis and the unity of biochemistry. *Cell (Cambridge, Mass.)* 72, 819-822.
- White, H. B. (1976). Coenzymes as fossils of an earlier metabolic state. *J. Mol. Evol.* 7, 101-104.
- White, R. H. (1980). Stoichiometry and stereochemistry of deuterium incorporated into fatty acids by cells of *Escherichia coli* grown on [methyl-²H₃]acetate. *Biochemistry* 19, 9-15.
- White, R. H. (1987). Biosynthesis of the 1,3,4,6-hexanetetra-carboxylic acid subunit of methanofuran. *Biochemistry* 26, 3161-3167.
- White, R. H. (1989a). A novel biosynthesis of medium chainlength γ -ketodicarboxylic acids in methanogenic archaeobacteria. *Arch. Biochem. Biophys.* 270, 691-697.
- White, R. H. (1989b). Biosynthesis of the 7-mercaptoheptanoic acid subunit of component B (7-mercaptoheptanoylthreonine phosphate) of methanogenic bacteria. *Biochemistry* 28, 860-865.
- White, R. H. (1989c). Steps in the conversion of α -ketosuberate to 7-mercaptoheptanoic acid in methanogenic bacteria. *Biochemistry* 28, 9417-9423.
- White, R. H. (1990). Biosynthesis of methanopterin. *Biochemistry* 29, 5397-5404.
- White, R. H. (1993a). Structure of the modified folates in the thermophilic archaeobacteria *Pyrococcus furiosus*. *Biochemistry* 32, 745-753.
- White, R. H. (1993b). Structures of the modified folates in the extremely thermophilic archaeobacteria *Thermococcus litoralis*. *J. Bacteriol.* 175, 3661-3663.
- White, R. H. (1994). Biosynthesis of 7-mercaptoheptanoylthreonine phosphate. *Biochemistry* 33, 7077-7081.
- White, R. H. (1996a). Biosynthesis of methanopterin. *Biochemistry* 35, 3447-3456.
- White, R. H. (1996b). Absolute stereochemistry of methanopterin and 5,6,7,8-tetrahydromethanopterin. *Chirality* 8, 332-340.
- White, R. H. (1997). Structural characterization of the modified folates in archaea. In "Methods in Enzymology" (D. B. McCormick, J. W. Suttie, and C. Wagner, eds.), Vol. 281, pp. 391-401. Academic Press, San Diego, CA.
- White, R. H. (1998). Methanopterin biosynthesis: Methylation of the biosynthetic intermediates. *Biochim. Biophys. Acta* 1380, 257-267.

- nd Hedderich, R. (1998). Two malate dehydrogenases from *Methanohalobium*. *Arch. Microbiol.* **170**, 38–42.
- 34). Expression and functional properties of the *mdhA* gene from *Methanohalobium*. *J. Biol. Chem.* **273**, 331–337.
- 397). A selenium-dependent and a selenindependent hydrogenase and their transcriptional regulation in *Methanohalobium*. *Mol. Microbiol.* **23**, 1033–1043.
- tion and properties of L-citramalate hydratase from *Methanohalobium*. *Biochemistry* **36**, 1033–1038.
- ito, H., Asahi, S., Nishimura, S., Katzev, J. A., and Okada, N. (1997). Biosynthesis of 7-deazaguanosine specific to archaeal base replacement on the tRNA polynucleotide chain. *Nucleic Acids Res.* **25**, 2015–2021.
- rrier protein reductases from *Escherichia coli*. In: *The Enzymes* (P. D. Boyer, ed.), Vol. 14, pp. 66–73. Academic Press, New York.
- ogenesis and the unity of biochemistry. *Science* **237**, 122–127.
- i earlier metabolic state. *J. Mol. Evol.* **7**, 1–10.
- hemistry of deuterium incorporated into nucleic acids by *Methanohalobium*. *Biochemistry* **36**, 1033–1038.
- n hexanetetra-carboxylic acid subunit of the *mdhA* gene product. *Biochemistry* **36**, 1033–1038.
- medium chainlength γ -ketodicarboxylic acids. *Biochem. Biophys. Res. Commun.* **270**, 691–697.
- aptoheptanoic acid subunit of component C of methanogenic bacteria. *Biochemistry* **36**, 1033–1038.
- f α -ketosubstrate to 7-mercaptoheptanoic acid. *Biochemistry* **36**, 1033–1038.
- erin. *Biochemistry* **29**, 5397–5404.
- folates in the thermophilic archaeobacteria. *Biochemistry* **35**, 3447–3456.
- folates in the extremely thermophilic archaeobacterium. *Biochemistry* **35**, 3447–3456.
- p-toheptanoylthreonine phosphate. *Biochemistry* **35**, 3447–3456.
- terin. *Biochemistry* **35**, 3447–3456.
- i of methanopterin and 5,6,7,8-tetrahydromethanopterin. *Biochemistry* **35**, 3447–3456.
- n of the modified folates in archaea. In: *The Enzymes* (J. W. Suttie, and C. Wagner, eds.), Vol. 18, pp. 1–10. Academic Press, San Diego, CA.
- s: Methylation of the biosynthetic intermediates. *Biochemistry* **35**, 3447–3456.
- White, R. H., and Zhou, D. (1993). Biosynthesis of coenzymes in methanogens. In: *The Enzymes* (J. C. Ferry, ed.), pp. 409–444. Chapman & Hall, New York.
- White, R. H., Roberts, M. F., and Gorkovenko, A. (1996). Naturally occurring 1,3,4,6-hexanetetra-carboxylic acid has the mesostereochemistry. *Chirality* **8**, 291–293.
- Worrell, V.F., and Nagle, D. P., Jr. (1988). Folic acid and pteroylpolyglutamate contents of archaebacteria. *J. Bacteriol.* **170**, 442–4420.
- Xiao, B., Shi, G. B., Chen, X., Yan, H. G., and Ji, X. H. (1999). Crystal structure of 6-hydroxymethyl-7,8-dihydropterin pyrophosphokinase, a potential target for the development of novel antimicrobial agents. *Struct. Folding Des.* **7**, 489–496.
- Xu, H., Aurou, R., Rose, G. D., and White, R. H. (1999). Excavating two ancient enzymes in archaea. *Nat. Struct. Biol.* **6**, 750–754.
- Yoon, K.-S., Ishii, M., Igarashi, Y., and Kodama, T. (1996). Purification and characterization of 2-oxoglutarate: Ferredoxin oxidoreductase from a thermophilic, obligately chemolithoautotrophic bacterium, *Hydrogenobacter thermophilus* TK-6. *J. Bacteriol.* **178**, 3365–3368.

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

☒ **BLACK BORDERS**

☒ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**

☐ **FADED TEXT OR DRAWING**

☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**

☐ **SKEWED/SLANTED IMAGES**

☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**

☐ **GRAY SCALE DOCUMENTS**

☒ **LINES OR MARKS ON ORIGINAL DOCUMENT**

☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**

☐ **OTHER:** _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.